

REVIEW

Dynamics of DNA damage response proteins at DNA breaks: a focus on protein modifications

Sophie E. Polo¹ and Stephen P. Jackson^{1,2}

The Gurdon Institute, Department of Biochemistry, University of Cambridge, Cambridge CB2 1QN, United Kingdom

Genome integrity is constantly monitored by sophisticated cellular networks, collectively termed the DNA damage response (DDR). A common feature of DDR proteins is their mobilization in response to genotoxic stress. Here, we outline how the development of various complementary methodologies has provided valuable insights into the spatiotemporal dynamics of DDR protein assembly/disassembly at sites of DNA strand breaks in eukaryotic cells. Considerable advances have also been made in understanding the underlying molecular mechanisms for these events, with post-translational modifications of DDR factors being shown to play prominent roles in controlling the formation of foci in response to DNA-damaging agents. We review these regulatory mechanisms and discuss their biological significance to the DDR.

Genome integrity is continuously challenged by DNA lesions, many thousands of which arise in each human cell every day (Lindahl and Barnes 2000). While the majority of these lesions occur as byproducts of normal cell metabolism or DNA replication, they are also induced by radiation and toxic environmental chemicals (Friedberg et al. 2006; Jackson and Bartek 2009; Ciccia and Elledge 2010). Programmed DNA lesions also form as intermediates during developmentally regulated genome rearrangements in lymphocytes and germ cells (Jackson and Bartek 2009; Longhese et al. 2009; Tsai and Lieber 2010). DNA damage can have deleterious effects, as it interferes with DNA replication and transcription, and because it can ultimately result in mutations and chromosomal aberrations. Genome integrity is preserved by DNA damage signaling and repair machineries, which counteract the adverse consequences of DNA lesions and prevent their transmission to daughter cells (Hoeijmakers 2001; Ciccia and Elledge 2010). Defects in DNA damage signaling or repair contribute to aging and various disorders, including developmental defects, neurodegenerative diseases, and cancer (Jackson and Bartek 2009), which highlights the

critical importance of an efficient DNA damage response (DDR) for cell and organism viability.

This review focuses on cellular responses to DNA single-strand breaks (SSBs) and double-strand breaks (DSBs) in eukaryotic cells, with an emphasis on mammalian systems. After a general overview of how such lesions are generated, signaled, and repaired, we outline recent work that has characterized the spatiotemporal dynamics of protein assembly/disassembly at sites of DNA breaks and the regulatory mechanisms involved. In particular, we highlight the major contribution of post-translational modifications in mediating and controlling these processes, and discuss the biological significance of DNA damage foci.

Cellular responses to DNA strand breaks

Formation and repair of DNA breaks

DNA DSBs are the most deleterious form of DNA damage because they do not leave an intact complementary strand to be used as a template for DNA repair. If left unrepaired, they can ultimately lead to chromosome breaks and translocations that are associated with developmental defects, neurodegeneration, immunodeficiency, radiosensitivity, sterility, and cancer predisposition (Jackson and Bartek 2009). DSBs are generated in response to ionizing radiation (IR) or radiomimetic drugs by free radical attack of deoxyribose, and also arise in cells treated with topoisomerase II inhibitors that prevent religation of DNA strands broken by topoisomerase II activity. DSBs can also form upon replication of DNA molecules containing other DNA lesions, such as DNA SSBs, and are produced by specific nucleases during V(D)J and class switch recombination in vertebrate lymphocytes, meiotic recombination in germ cells, mating type switching in yeast, apoptotic cell death, and retroviral integration (Friedberg et al. 2006; Jackson and Bartek 2009). Last, but not least, naturally occurring DSBs at chromosome ends are associated with human cell aging, as they are exposed when telomeres become critically short during replicative senescence (d'Adda di Fagagna et al. 2003).

Cells rely on two major pathways to repair DSBs: homologous recombination (HR) and nonhomologous

[*Keywords:* checkpoint; DNA breaks; DNA damage foci; post-translational modifications; repair]

¹The authors contributed equally to this work.

²Corresponding author.

E-MAIL s.jackson@gurdon.cam.ac.uk; FAX 44-1223-334-089.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.2021311>.

Polo and Jackson

end-joining (NHEJ) (Hartlerode and Scully 2009; Pardo et al. 2009). These pathways are complementary and operate optimally under different circumstances. HR requires the presence of a homologous template, usually a sister chromatid, which allows accurate repair of post-replicative DSBs in S and G2 phases of the cell cycle (San Filippo et al. 2008; Moynahan and Jasin 2010). In contrast, NHEJ can operate throughout the cell cycle without the need for template DNA, and is often mutagenic because deletions or insertions can be induced at sites of repair (Lieber 2010). NHEJ, which is the prevalent DSB repair pathway in higher eukaryotes, essentially mediates the direct ligation of broken DNA ends, and usually involves minimal DNA end processing. In NHEJ, DNA ends are first bound by the Ku70/Ku80 heterodimer, which recruits and activates the DNA-dependent protein kinase catalytic subunit (DNA-PKcs) to form the DNA-PK holoenzyme (Gottlieb and Jackson 1993). Broken DNA ends juxtaposed by DNA-PK are then acted on by factors such as the nuclease Artemis, polynucleotide kinase (PNK), Aprataxin, and APLF (Aprataxin and PNK-like factor) before being ligated by the XLF-XRCC4 (X-ray cross-complementing-4)-LigaseIV complex (Table 1; Lieber and Wilson 2010). In addition, alternative end-joining pathways sometimes operate, particularly when classical

NHEJ is impeded because some NHEJ component is missing or mutated. These alternative end-joining pathways often rely on terminal microhomologies for the joining reaction, and involve certain factors that also function in HR or SSB repair (SSBR), such as the MRE11-RAD50-NBS1 (MRN) complex, poly(ADP-ribose) polymerase-1 (PARP-1), XRCC1, and DNA Ligase I or III (McVey and Lee 2008).

An important regulatory step that determines the choice between the two main DSB repair pathways is the process of DSB resection, which is required for HR but not NHEJ. Resection comprises the 5'-to-3' nucleolytic processing of DNA ends by the MRN complex (Rupnik et al. 2010) in conjunction with auxiliary factors including CtIP, RECQ family helicases, and the nucleases Exo1 and Dna2 (Zou and Elledge 2003; Bernstein and Rothstein 2009; Mimitou and Symington 2009; Huertas 2010; Longhese et al. 2010; You and Bailis 2010). Resulting ssDNA overhangs are then coated by the ssDNA-binding complex RPA (replication protein A) before being substituted by RAD51 proteins with the help of factors such as RAD51 paralogs, RAD52, and other proteins that comprise the FA (Fanconi anemia) pathway such as FANCD1 (FA-associated nuclease CD1)/BRCA2 (breast cancer-2, early onset) and FANCN/PALB2 (partner and localizer of

Table 1. Factors involved in DNA strand break repair and damage signaling in budding yeast and mammals

	Mammals	Yeast (<i>S. cerevisiae</i>)
<u>DNA strand break repair</u>		
NHEJ		
End binding	MRE11-RAD50-NBS1(MRN) Yku70-Yku80 DNA-PKcs	Mre11-Rad50-Xrs2 (MRX) Ku70-Ku80
End processing	Artemis, APLF, PNK, APTX	
Ligation	LigaseIV-XRCC4-XLF	Lig4-Lif1-Nej1
HR		
Resection	MRN, CtIP, EXO1, BLM, DNA2?	MRX, Sae2, Exo1, Sgs1, Dna2
Homologous pairing and strand exchange	RPA, RAD51, RAD52, RAD54, RAD51 paralogs BRCA2-PALB2 (=FANCD1-FANCN)	Rfa, Rad51, Rad52, Rad54, Rad55-Rad57
DNA synthesis	PCNA, Pol δ	PCNA, Pol δ
HR resolvases	MUS81-EME1, GEN1, SLX1-SLX4, XPF-ERCC1	Mus81-Eme1, Yen1 Slx1Slx4, Rad1-Rad10
Dissolution of HR intermediates	BLM-TOPOIII-RMI1-RMI2, RTEL1	Sgs1-Top3-Rmi1, Srs2
SSBR		
Detection	PARP-1	
End processing	APE1, XRCC1, PNK, APTX	
Gap filling, ligation	LigIII, Pol β	
<u>DNA damage signaling</u>		
Sensors		
	MRN RPA (+RFC-like, PCNA-like checkpoint clamp)	MRX Rfa (+RFC-like, PCNA-like checkpoint clamp)
Transducers		
	ATM ATR-ATRIP	Tel1 Mec1-Ddc2
Mediators		
ATM signaling	53BP1, MDC1, BRCA1, MCPHI PTIP	Rad9
ATR signaling	TopBP1 Claspin	Dpb11 Mrc1
Effectors		
	CHK1 CHK2	Chk1 Rad53

BRCA2) that play key roles in detecting and repairing interstrand cross-links, particularly at sites of stalled DNA replication (Moldovan and D'Andrea 2009). The RAD51 nucleofilament, together with various other HR factors, then mediates homology search in the sister chromatid, followed by strand invasion into the homologous template. After the actions of DNA polymerases and DNA end ligation by Ligase I, DNA helicase and resolvase enzymes then mediate the cleavage and resolution of HR intermediates to yield intact, repaired DNA molecules (Table 1; Mazon et al. 2010).

Although not as harmful as DSBs, SSBs are toxic to the cell, as they can block DNA replication and transcription, and, indeed, SSB defects are associated with several hereditary neurodegenerative diseases (Caldecott 2008). SSBs form upon oxidative attack of deoxyribose by free radicals arising from cell metabolism or through exposure to agents such as H₂O₂, IR, and radiomimetic drugs. They also arise as intermediates during excision repair of base damage, and upon inhibition of topoisomerase I. In most eukaryotes, SSBs are initially detected by PARP-1, whose binding to DNA breaks triggers poly-(ADP-ribosylation) of numerous nuclear proteins, including itself. These modifications in turn promote the binding of XRCC1, which acts as a molecular scaffold for downstream SSB components involved in end-processing, gap filling, and ligation (Table 1).

Signaling of DNA breaks

DNA repair is tightly coordinated with cell cycle progression through the activation of orchestrated signaling pathways that are often termed DNA damage checkpoints (Harrison and Haber 2006; Harper and Elledge 2007; Lazzaro et al. 2009). In response to unrepaired DNA damage, these pathways delay or stop the cell cycle at critical stages before or during DNA replication (G1/S and intra-S checkpoints) and before cell division (G2/M checkpoint), thereby preventing duplication and segregation of damaged DNA. DNA damage signaling cascades are complex, coordinated events that require the actions of various proteins whose functions can be categorized as DNA damage sensors, transducers, mediators, and effectors. Of particular note are the MRN sensor complex that detects DSBs (Lavin 2007), and RPA that signals the accumulation of ssDNA resulting from DNA damage processing. MRN contributes to the recruitment and activation of the apical DDR kinase ATM (ataxia telangiectasia mutated) (Uziel et al. 2003; Falck et al. 2005; Lee and Paull 2005), while RPA recruits the ATR (ATM and rad3-related) kinase via its partner protein, ATRIP (ATR-interacting partner) (Cortez et al. 2001; Rouse and Jackson 2002; Zou and Elledge 2003). With the help of mediator proteins—such as MDC1 (mediator of DNA damage checkpoint), 53BP1 (p53-binding protein 1), and BRCA1 for ATM, and TopBP1 (topoisomerase-binding protein 1) and Claspin for ATR—transducer kinases activate the effector kinases Chk1 and Chk2, which then spread the signal throughout the nucleus. Ultimate targets of these signaling cascades include transcription factors, cell cycle

regulators, the apoptotic machinery, and DNA repair factors (Harrison and Haber 2006; Harper and Elledge 2007; Lazzaro et al. 2009).

Responses to DNA breaks in a physiological context

Signaling and repair of DNA breaks occur in a context where they can potentially interfere with other DNA metabolic activities such as replication and transcription. Indeed, ATM has been reported to mediate local inhibition of both RNA polymerase I- and II-dependent transcription at sites of DNA breaks in human cells (Kruhlak et al. 2007; Shanbhag et al. 2010). Such transcriptional inhibition is accompanied by the clearance of active RNA polymerases from damage sites, highlighting that protein dynamics at DNA breaks operate in both directions: While DNA damage signaling and repair factors are recruited to DNA breaks, other factors dissociate (Fig. 1).

When studying the DDR in its cellular context, one also has to consider that the physiological substrate for the DDR machinery in the cell nucleus is DNA wrapped around histone proteins in the form of chromatin, the basic unit of which is the nucleosome (Kornberg 1977; Luger et al. 1997). Nucleosomes can be further compacted into higher-order chromatin structures with linker histones and nonhistone components such as heterochromatin protein 1 (HP1) (McBryant et al. 2006; Fanti and Pimpinelli 2008). Although chromatin acts as a physical barrier to the detection and repair of DNA lesions, it is also a dynamic structure that can be modulated by DNA methylation (Kulis and Esteller 2010), incorporation of histone variants (Bernstein and Hake 2006), histone post-translational modifications (Kouzarides 2007), and nucleosome repositioning by ATP-dependent remodeling complexes (Clapier and Cairns 2009). Moreover, a growing body of evidence highlights the importance of such modulations of chromatin organization in the DDR (for review, see Misteli and Soutoglou 2009; van Attikum and Gasser 2009), the most prominent and widely documented being phosphorylation of the histone variant H2AX (γ H2AX) identified >10 years ago (Rogakou et al. 1998). As a consequence, and as described in further detail below, it comes as no surprise that chromatin-associated proteins are also mobilized to and from DNA breaks (Fig. 1; Table 2). In particular, chromatin remodeling complexes and histone chaperones have been linked to histone mobilization (Tsukuda et al. 2005; Kent et al. 2007; Shim et al. 2007; Xu et al. 2010), histone variant replacement (Kusch et al. 2004; Papamichos-Chronakis et al. 2006; Heo et al. 2008), and new histone deposition (Polo et al. 2006; Chen et al. 2008) at sites of DNA breaks.

Methods for studying DDR protein assembly/disassembly at DNA breaks

Our understanding of DDR protein dynamics at DNA breaks has been greatly advanced through the development of various methodologies for studying DDR protein assembly and disassembly at damage sites *in vitro* and *in*




Figure 1. Protein dynamics to and from sites of DNA breaks. DNA damage checkpoint and repair factors and modulators of chromatin organization are recruited (green arrows) to DNA breaks (SSB and DSB), while transcription machineries are excluded from DDR foci (red arrows), and the dynamics of structural chromatin components operate in both directions (orange arrows).

vivo (summarized in Fig. 2; Lukas et al. 2005; Nagy and Soutoglou 2009). Initially, the recruitment of DDR proteins to broken DNA was analyzed in vitro by biochemical assays employing various types of DNA substrates (e.g., circular vs. linear DNA), as described for the NHEJ factors Ku and DNA-PKcs, which display strong affinity for DNA ends (Gottlieb and Jackson 1993; Dynan and Yoo 1998). More recently, advanced biophysical approaches have enabled single-molecule imaging of DDR proteins such as yeast Rad54 and human RAD51 on DNA (Amitani et al. 2010).

In vivo studies of DDR protein recruitments include imaging of protein dynamics at sites of DNA breaks and biochemical approaches to analyze changes in protein binding to damaged chromatin. When DNA damage is inflicted on the whole nucleus by exposing cells to genotoxic agents such as H_2O_2 , IR, radiomimetic drugs or topoisomerase inhibitors, protein accumulation at DNA damage sites can take the form of cytologically discernable foci, as initially observed for the proteins Rad51, BRCA1, and MRN in human cells (Haaf et al. 1995; Maser et al. 1997; Scully et al. 1997). Such foci can be visualized by indirect immunofluorescence or in real time by imaging live cells expressing fluorescently tagged proteins. Importantly, the number of focus-positive cells and the number of foci per cell each increase in a dose-dependent manner in mammalian cells (for example, 20–40 DSB foci per nucleus per Gray of radiation for a mammalian cell), and focus number and size also change over time after DNA damage induction (van Veelen et al. 2005). In this regard, it is noteworthy that similar foci can be detected at recombination sites in meiotic cells, as initially illustrated by the local accumulation of the RAD51 recombinase in vertebrate germ cells (Ashley et al. 1995). Another approach for analyzing protein dynamics in response to DNA breaks consists of monitoring increased chromatin binding of proteins

in damaged versus undamaged cells by biochemical cell fractionation (Drouet et al. 2005), a method that was converted recently into high-throughput formats by the implementation of SILAC (stable isotope labeling by amino acids in cell culture)-based mass spectrometry (Chou et al. 2010; Larsen et al. 2010).

The recruitment of DDR factors to DNA breaks can also be analyzed by chromatin immunoprecipitation (ChIP) or cell imaging upon local induction of DNA damage by microirradiation or specific endonucleases. The generation of sequence-specific DSBs by selected endonucleases was pioneered with yeast homothallic (HO) endonuclease (Sugawara and Haber 2006), translated into mammalian cells by using homing endonucleases such as I-SceI (Jasin 1996) and I-PpoI (Berkovich et al. 2008), and more recently developed into genome-wide formats by combining expression of the restriction enzyme AsiSI with microarray hybridization and high-throughput sequencing (Iacovoni et al. 2010; Massip et al. 2010). Furthermore, generation of DSBs by the targeting of a nuclease to a defined locus in the genome can be attained by fusing its nuclease domain to a zinc finger protein or a Lac repressor, as was done recently with the FokI enzyme (Urnov et al. 2005; Shanbhag et al. 2010).

In addition to the use of endonucleases, several localized irradiation techniques have been developed to introduce discrete sites of DNA breaks in the cell nucleus, including partial cell volume irradiation with ultrasoft X-rays through a gridded shield (Nelms et al. 1998) or high-voltage X-rays through a microcollimator (Pataky et al. 2009). Similarly, local UV-C irradiation through micropore filters combined with the expression of UVDE (UV-specific endonuclease) in cells defective in repairing UV lesions has been used to generate localized SSBs (Okano et al. 2003). DNA breaks, together with various base alterations, can also be induced locally by laser microirradiation, with or

Table 2. Factors involved in chromatin dynamics recruited to/dissociating from damaged chromatin in response to DNA breaks

	Name	Organism	References
<u>DNA methyltransferases</u>			
	Dnmt1	Human, mouse	Mortusewicz et al. 2005; Cuozzo et al. 2007; O'Hagan et al. 2008
	Dnmt3b	Human	O'Hagan et al. 2008
<u>Histone-modifying enzymes</u>			
Histone methyltransferase	EZH2	Human	O'Hagan et al. 2008; Chou et al. 2010
	PR-Set7/Set8	Human	Oda et al. 2010
Histone acetyltransferase	Esa1	Budding yeast	Tamburini and Tyler 2005
	Gcn5	Budding yeast	Tamburini and Tyler 2005
	Hat1	Budding yeast	Qin and Parthun 2006
	NuA4	Budding yeast	Downs et al. 2004
	Tip60	Human	Murr et al. 2006
	Histone deacetylase	Hst1	Budding yeast
Rpd3		Budding yeast	Tamburini and Tyler 2005
Sir2		Budding yeast	Tamburini and Tyler 2005
HDAC1		Human	Miller et al. 2010; Polo et al. 2010
HDAC2		Human	Miller et al. 2010
HDAC4		Human	Kao et al. 2003
SIRT1		Human, mouse	O'Hagan et al. 2008; Oberdoerffer et al. 2008
SIRT6		Human	Kaidi et al. 2010
<u>Chromatin remodeling factors</u>			
	INO80	Budding yeast	Morrison et al. 2004; van Attikum et al. 2004
	RSC	Budding yeast	Chai et al. 2005; Shim et al. 2005
	SWI/SNF	Budding yeast	Chai et al. 2005
	SWR1	Budding yeast	van Attikum et al. 2007
	ALC1	Human	Ahel et al. 2009
	INO80	Human, mouse	Kashiwaba et al. 2010
	ISWI	Human	Erdel et al. 2010
	NuRD	Human	Chou et al. 2010; Larsen et al. 2010; Polo et al. 2010; Smeenk et al. 2010
	p400	Human	Xu et al. 2010
	SWI/SNF	Human	Park et al. 2006; Peng et al. 2009; Lee et al. 2010
<u>Histone chaperones</u>			
	CAF-1	Human	Moggs et al. 2000; Okano et al. 2003; Polo et al. 2006
	FACT	Human	Huang et al. 2006; Heo et al. 2008

All listed factors are recruited to damaged chromatin, with the exception of FACT, which dissociates from chromatin upon DNA damage.

without cell photosensitization with halogenated thymidine analogs such as bromodeoxyuridine (BrdU) or a DNA intercalating agent like Hoechst (Limoli and Ward 1993; Lukas et al. 2005). This technique has also been refined recently by the use of highly focused multiphoton lasers

(Mari et al. 2006; Botchway et al. 2010). In such studies, it is important to calibrate the laser wavelength, energy output, and photosensitization method used, as these determine the type and density of DNA lesions. Other techniques such as localized cell irradiation with heavy-ion or




Figure 2. Methods for studying the recruitment of DDR proteins to DNA breaks. Scheme describing the multiple methods used to generate or mimic DNA breaks in vitro and in vivo and the techniques employed to monitor recruitment of DDR factors to such breaks.

α -particle microbeams (Jakob et al. 2005; Stap et al. 2008) elicit similar tracks of DNA breaks and base modifications in the cell nucleus. Other ways of triggering DNA DSB signaling in mammalian cells without the need for any exogenous DNA-damaging agent are telomere shortening in senescent cells and artificial telomere uncapping by inducible loss of function of a component of the Shelterin complex that protects chromosome ends, both of which trigger the formation of DDR foci at telomeres (d'Adda di Fagagna et al. 2003; Takai et al. 2003). Similar uncapping strategies have been used in yeast (Wellinger 2010). Complementary to the aforementioned methods, photobleaching techniques such as fluorescence recovery after photobleaching (FRAP) have provided valuable insights into the dynamics of DDR proteins by measuring their turnover rates at damage sites (Essers et al. 2006; Mortusewicz et al. 2008), revealing that DDR factors are not stably bound within a DDR focus but dynamically exchange with neighboring nuclear compartments. Collectively, these studies have proven critical for understanding the hierarchy and spatiotemporal dynamics of DDR factor accumulation at DNA breaks.

Spatiotemporal dynamics of DDR protein assembly/disassembly at DNA breaks

To ensure the efficient signaling and repair of DNA damage, DDR proteins must relocate to the right place at the right time. As described below, a wealth of studies have highlighted the tight spatiotemporal coordination of DDR protein assembly and disassembly at DNA breaks.

Spatial organization of DDR foci

One hallmark of DDR proteins is their local accumulation at damage sites, which in many cases leads to the formation of discrete, cytologically detectable foci. Notably, however, not all DDR factors accumulate at DNA breaks in a manner that can be readily observed under the microscope. This is exemplified by NHEJ components, as described below, and the Chk1 and Chk2 effector kinases, which become phosphorylated at damage sites but then quickly dissociate and distribute throughout the nucleus (Lukas et al. 2003; Smits et al. 2006). Furthermore, factors accumulating at DSB sites do not always colocalize perfectly, with two classes of DDR proteins being readily distinguishable based on their spatial distributions (Fig. 3): those present directly at damage sites, coating ssDNA resulting from DSB resection, and those associated with DSB-flanking chromatin (Bekker-Jensen et al. 2006). In this regard, we point out that, because NHEJ components are usually confined close to the DSBs themselves and do not spread substantially into adjacent chromatin, such factors are not usually evident at radiation-induced or even laser-induced foci unless high levels of damage are used (Bekker-Jensen et al. 2006). However, they can be readily detected at DSB sites by ChIP (Zhang et al. 2007; Miller et al. 2010) or cellular fractionation methods (Drouet et al. 2005).

Strikingly, for various DSB-associated DDR factors and protein modifications, their foci increase in size over

time, which reflects these factors and modifications spreading away from the DSB into adjacent chromatin (Fig. 3), as initially observed for γ H2AX in mammalian cells (Rogakou et al. 1999). γ H2AX spreading was also described in yeast (Downs et al. 2004; Shroff et al. 2004), although over shorter distances: up to 100 kb as opposed to several megabases in mammals (Rogakou et al. 1999; Meier et al. 2007; Iacovoni et al. 2010). Another prominent difference reported between yeast and higher eukaryotes lies in the mobility of DDR foci. In budding yeast, DDR foci are highly mobile and coalesce into repair centers (Lisby et al. 2003), which contrasts with the reported relative positional stability of broken DNA ends in mammalian cells (Nelms et al. 1998; Kruhlak et al. 2006; Soutoglou et al. 2007; Jakob et al. 2009). The reason for this apparent discrepancy is not yet clear, and might simply relate to the difference in nuclear size between these organisms. For example, it is possible that mammalian DDR foci are able to move distances similar to those traversed by foci in yeast cells, but this might not have been evident in the mammalian studies because such movements would occur only within a small fraction of the total nuclear volume. Indeed, recent studies have indicated that some DSB mobility does occur in mammalian cells to promote joining of distal DNA breaks (Difilippantonio et al. 2008; Dimitrova et al. 2008).

The dynamics of DDR foci are also dictated by chromatin organization and transcriptional activity, as they clearly differ between relaxed, highly transcribed euchromatin




Figure 3. Spatial organization of DDR protein accumulation at DNA DSBs. (A) DDR signal spreading. DDR proteins initially accumulate at DSB sites and then spread at distance via a positive feedback loop involving MDC1, which binds γ H2AX, the MRN complex, and ATM kinase, which phosphorylates additional H2AX molecules further away from the break site. (B) Regional distribution of DDR proteins around DSBs. Factors involved in ATR signaling accumulate proximal to the break site on ssDNA generated by DNA end resection, while ATM signaling factors localize on flanking chromatin regions.

and more compact heterochromatin that concentrates in specific chromosomal domains such as centromeres and telomeres. For example, it has been shown that γ H2AX does not propagate effectively on actively transcribed genes in human cells (Iacovoni et al. 2010). Furthermore, despite the fact that DNA breaks can be efficiently induced in heterochromatin, this chromatin compartment is generally refractory to γ H2AX focus formation in yeast and mammalian cells (Cowell et al. 2007; JA Kim et al. 2007). This barrier is relieved in S phase, however, by DNA replication, and replication stress-associated γ H2AX is also detected in heterochromatin regions (Chadwick and Lane 2005; Cowell et al. 2007; Rozenzhak et al. 2010; Szilard et al. 2010), suggesting that transient decompaction of heterochromatin and/or displacement of heterochromatin components upon DNA replication is required for the propagation of H2AX phosphorylation. Notably, the γ H2AX foci that persist longest after damage in mammalian cells generally localize at the periphery of heterochromatin domains, suggesting that heterochromatic DSBs are refractory to repair; indeed, it has been shown that this γ H2AX persistence can be alleviated by depletion of heterochromatin components (Goodarzi et al. 2008).

Temporal organization of DDR foci

In addition to shaping our knowledge of DDR protein distributions in response to DNA breaks, live-cell imaging by time-lapse microscopy has substantially contributed to our understanding of DDR protein dynamics in both yeast and mammalian cells, especially in regard to the early kinetics of DDR focus formation. Furthermore, the use of siRNA-mediated depletion methods or mutant cell lines has been instrumental in functionally dissecting the sequential recruitment of DDR proteins to DNA breaks (e.g., Lisby et al. 2004; Mari et al. 2006; Mailand et al. 2007). Indeed, multiple lines of evidence support the idea that DDR proteins assemble in a sequential, coordinated manner at sites of DNA breaks, rather than being recruited as a preformed protein complex (Fig. 4A). For instance, the recruitment of DDR factors to SSBs is very rapid and transient (Mortusewicz et al. 2007), reaching maximal accumulation within 1–2 min, due to its dependency on poly(ADP-ribosylation) (PARylation), which is a relatively short-lived protein modification (Gagne et al. 2006; Hakme et al. 2008). Factors involved in DSB repair by NHEJ are also recruited within seconds upon break formation and normally dissociate within 2 h, while HR factors show delayed and persistent recruitment to DSBs, reflecting different repair kinetics between these two pathways (Kim et al. 2005; Mari et al. 2006; Mailand et al. 2007; Uematsu et al. 2007; Yano et al. 2008; Doil et al. 2009).

The accumulation kinetics of factors involved in DSB signaling has also been studied in great detail, and has been described as a two-stage process in which initial recruitment occurs independently of H2AX phosphorylation, followed by sustained DDR factor retention in a γ H2AX-dependent manner (Celeste et al. 2003b; Yuan et al. 2010). As discussed in the following sections, studies from multiple laboratories have built up a fairly comprehensive




Figure 4. Temporal regulation of DDR protein accumulation at DNA breaks. (A) Sequential recruitment of DDR factors to SSBs and DSBs generated by laser microirradiation. (B) Cell cycle regulation of DDR foci formation. (Solid line) Efficient focus formation; (dashed line) weak/undetected foci.

picture of the mechanisms involved in γ H2AX-dependent focus formation. In contrast, our knowledge of the initial γ H2AX-independent recruitment of DSB signaling factors is limited, although it seems to involve the MRN complex, additional histone modifications, and changes in chromatin compaction at damage sites (FitzGerald et al. 2009; Xie et al. 2009; Yuan and Chen 2010).

Importantly, responses to DSBs can be markedly influenced by cell cycle status (Fig. 4B). Thus, while focal accumulation of DDR factors such as γ H2AX, MRN, and MDC1 occurs regardless of the cell cycle stage, focus formation by others—including CtIP, RPA, ATRIP-ATR, BRCA1, and RAD51—takes place effectively only in association with ssDNA formation by DNA end resection in S/G2 cells (Lisby et al. 2004; Bekker-Jensen et al. 2006; Jazayeri et al. 2006; Sartori et al. 2007). In contrast, resection has been reported in G1 at sites of IR-induced breaks in budding yeast, although the rate of resection is higher in S phase and lesions appear to become recognized by the HR machinery only upon entry into S phase (Barlow et al. 2008). Recent work has shown that cell cycle control of DNA resection and HR relies on cell cycle-regulated expression of key factors such as CtIP and BRCA1, and is also governed by cyclin-dependent kinases (Cdks) that phosphorylate evolutionarily conserved residues in CtIP (Huertas et al. 2008; Huertas and Jackson 2009; You and Bailis 2010).

In addition to regulation during interphase, recent observations have shed light on a distinct response to DSBs during mitosis, with only partial activation of the DDR

Polo and Jackson

taking place in this cell cycle stage (Fig. 4B). While the molecular basis of this DDR abrogation is still elusive, it appears to operate at the level of ubiquitylating enzymes that control 53BP1 and BRCA1 focus formation (Nelson et al. 2009; Giunta et al. 2010; Nakamura et al. 2010; van Vugt et al. 2010). It is speculated that early DNA damage signaling takes place in mitosis as a priming event for full DDR activation in the following G1 phase (Giunta et al. 2010).

Molecular mechanisms of DDR protein assembly/disassembly at DNA breaks

Analyzing the spatiotemporal dynamics of DDR proteins at DNA breaks has gone beyond being a descriptive approach, and has provided substantial mechanistic insights into how DDR factors assemble and disassemble at damage sites. In this section, we describe how DNA breaks are recognized by DNA damage sensors, how this drives focal recruitment of multiple downstream factors, and how protein modifications regulate these processes.

Direct recognition of DNA breaks by DDR factors

Among the first proteins recruited to DNA breaks are those able to directly recognize DNA breaks. Such factors bind broken DNA in a sequence-independent manner, and thus act as molecular sensors of DNA breaks. For example, PARP-1 and PARP-2 catalytic activity is triggered by them binding directly to SSBs and DSBs (Benjamin and Gill 1980; Ohgushi et al. 1980; de Murcia and Menissier de Murcia 1994; D'Amours et al. 1999). DNA binding is mediated via N-terminal zinc finger domains in PARP-1 (Menissier-de Murcia et al. 1989), and a basic N-terminal domain in PARP-2 (Ame et al. 1999). Similarly, the Ku70–Ku80 heterodimer and the MRN complex are DSB sensors that display direct binding to DNA ends *in vitro* (Mimori and Hardin 1986; de Jager et al. 2001), and are among the earliest factors to bind to DSBs *in vivo* (Lisby et al. 2004; Kim et al. 2005). Structural studies have shown that Ku70 and Ku80 form a ring-shaped heterodimer that encircles DNA and threads onto DNA by way of a DSB terminus (Walker et al. 2001), while the MRN complex binds DSBs via a globular head region comprised of MRE11 together with joined RAD50 ATPase domains, with MRE11 dimerization ensuring stable DNA binding, and RAD50 dimerization tethering DNA ends together (de Jager et al. 2001; Williams et al. 2008). As can be anticipated for factors binding the similar DNA structures, Ku competes with the budding yeast MRX complex (and, presumably, also with mammalian MRN) for binding to DSBs (Zhang et al. 2007; Clerici et al. 2008), and Ku also competes with PARP-1 for DNA end binding in mammalian cells (Wang et al. 2006).

Other DDR factors become recruited to processed DSBs via direct binding of evolutionarily conserved OB-fold motifs (Flynn and Zou 2010) to ssDNA generated by DNA end resection. The most extensively studied of these proteins is the ssDNA-binding complex RPA (Wold 1997), which accumulates at resected DSBs in an MRN- and CtIP-dependent manner. RPA in turn directs the

recruitment of the Rad9–Rad1–Hus1 (9-1-1) complex, a heterotrimeric ring that is structurally similar to the replicative sliding clamp PCNA (proliferating cell nuclear antigen) and is loaded onto ssDNA–dsDNA junctions by an RFC (replication factor C)-like clamp loader (Parrilla-Castellar et al. 2004). The ssDNA–dsDNA junction is also a substrate for binding by helicases and nucleases involved in DNA break repair (Mimitou and Symington 2009; Bernstein et al. 2010). Recently, two novel heterotrimeric ssDNA-binding complexes have been characterized, termed sensors of ssDNA 1 (SOSS1) and SOSS2, which contain the ssDNA-binding proteins hSSB1 and hSSB2, respectively (Richard et al. 2008; Huang et al. 2009a; Li et al. 2009; Skaar et al. 2009). Unlike RPA, these proteins appear to form DDR foci independently of cell cycle stage in a CtIP-independent manner, and recent work suggests that they could function upstream of MRN (Richard et al. 2010).

DDR protein assembly mediated by protein–protein interactions

Downstream from proteins that directly sense DNA breaks, other DNA damage signaling and repair factors are sequentially recruited to DNA lesions. The critical importance of protein–protein interactions in building such DDR factor assemblies is well exemplified by MDC1, which serves as a binding platform for DNA damage checkpoint and repair proteins (Fig. 5; Jungmichel and Stucki 2010). Similarly, XRCC1 and XRCC4 act as scaffold proteins, promoting the accumulation of repair factors at SSBs and DSBs, respectively (Fig. 5; Mortusewicz et al. 2008). The pivotal role of protein–protein interactions in the sequential association of repair factors to DDR foci is also illustrated by the recruitment of HR proteins to DSBs (Huen et al. 2010; Moynahan and Jasin 2010). For instance, in human cells, BRCA1 promotes the localization of BRCA2 to damage foci through the BRCA2-binding protein PALB2 (Xia et al. 2006; Sy et al. 2009; Zhang et al. 2009a,b). Through its interaction with RAD51, BRCA2 in turn promotes RAD51 assembly onto ssDNA (Jensen et al. 2010; J Liu et al. 2010; Thorslund et al. 2010).

In the NHEJ pathway, the Ku70–Ku80 heterodimer plays a central role in recruiting other NHEJ components by protein–protein interactions. In particular, Ku recruits DNA-PKcs (Dvir et al. 1992; Gottlieb and Jackson 1993) via a specific interaction between DNA-PKcs and the Ku80 C terminus (Gell and Jackson 1999; Singleton et al. 1999). Ku is also involved in recruiting the downstream NHEJ complex XLF–XRCC4–LigaseIV to DNA ends in mammalian cells (Nick McElhinny et al. 2000; Calsou et al. 2003; Yano et al. 2008), with analogous interactions also taking place in budding yeast (Teo and Jackson 2000; Palmos et al. 2008).

Regarding the apical checkpoint kinases, ATM is recruited to DSBs by the MRN/X complex (Uziel et al. 2003; Falck et al. 2005; Lee and Paull 2005) by a mechanism that involves direct and evolutionarily conserved interactions between ATM and the NBS1/Xrs2 C terminus, as shown in budding and fission yeast, *Xenopus*, and




Figure 5. Binding platforms at DNA breaks. NBS1, MDC1, XRCC1, and XRCC4 act as binding platforms for the recruitment of other DDR factors to DNA breaks promoting DNA damage signaling and/or repair. Dotted lines indicate protein-protein interactions, while horizontal lines at the end of the dotted lines indicate interacting regions. Some interactions involve post-translational modifications. (P) Phosphorylation; (PAR) PARylation. The red and the green semicircles represent BRCT and FHA domains, respectively. (b) Basic region at the end of the first BRCT domain of XRCC1 that interacts with poly[ADP-ribose]ylated PARPs.

human cells (Nakada et al. 2003; Falck et al. 2005; You et al. 2005). In contrast, ATR is recruited to resected DSBs via its interacting partner, ATRIP (Cortez et al. 2001), which binds to RPA-coated ssDNA in human cells, a mechanism that is conserved for yeast orthologs of these factors (Rouse and Jackson 2002; Zou and Elledge 2003; Ball et al. 2007). More generally, it appears that there is a conserved mechanism for apical DDR kinase recruitment to DNA damage sites that involves similar interaction motifs in the C termini of human Ku80-, NBS1-, and ATRIP-mediated interactions with DNA-PKcs, ATM, and ATR, respectively (Falck et al. 2005).

Control of DDR protein recruitment by post-translational modifications

As can be expected for a crucial cellular process, the building of multiprotein assemblies at DNA breaks is tightly controlled. Much of this is achieved by post-translational protein modifications (Fig. 6) that promote the recruitment or dissociation of DDR factors or regulate their residence times at damage sites. While such control mechanisms rely heavily on phosphorylation, a major breakthrough in recent years has been the realization that they are also regulated by other post-translational modifications, including ubiquitylation, sumoylation, methylation, acetylation, and PARylation.

Key roles for protein phosphorylation in the assembly of DDR foci

The central role of protein kinases and the widespread importance of phosphorylation and dephosphorylation reactions in the DDR are widely acknowledged, with several hundreds of phosphorylated targets having already been identified by mass spectrometry-based screens (Matsuoka et al. 2007; Bennetzen et al. 2010; Bensimon et al. 2010). While some of these phosphorylations likely serve to directly regulate the structure and activity of DDR target proteins, in many cases they act by providing regulated docking sites for other DDR factors. In this regard, DDR proteins frequently display phospho-binding motifs such as BRCT (breast cancer C-terminal) or FHA (Forkhead-associated) domains (Bork et al. 1997; Callebaut and Mornon 1997; Durocher et al. 1999; Mohammad and Yaffe 2009) that play pivotal roles in mediating the phospho-dependent assembly of DDR protein complexes (Fig. 6).

The prime example of a DDR kinase substrate is the histone variant H2AX, which is phosphorylated on a conserved C-terminal serine residue by ATM, ATR, and DNA-PK (Rogakou et al. 1998; Downs et al. 2000; Burma et al. 2001; Ward and Chen 2001; Stiff et al. 2004). This phosphorylation directs the assembly of downstream DDR components, including checkpoint mediators such as human MDC1 and MCPH1 (Microcephalin; also named BRIT1) and their budding and fission yeast orthologs, Rad9 and Crb2, respectively (Nakamura et al. 2004; Lee et al. 2005; Stucki et al. 2005; Hammet et al. 2007; Wood et al. 2007; Sanders et al. 2010; Sofueva et al. 2010). Phosphorylated H2AX also promotes the recruitment of chromatin-modifying complexes, including yeast NuA4, INO80, and SWR1; *Drosophila* Tip60; and human p400 (Downs et al. 2004; Kusch et al. 2004; Morrison et al. 2004; van Attikum et al. 2004, 2007; Xu et al. 2010). Most prominent among the various factors recruited to γ H2AX are MDC1 and its counterparts, which have been shown through structural and biochemical studies to bind directly to phospho-H2AX via their BRCT domains (Lee et al. 2005; Stucki et al. 2005; Hammet et al. 2007; Kilkenny et al. 2008). Notably, MDC1 binding to γ H2AX can also be modulated by neighboring modifications within the H2AX C-terminal tail, such as phosphorylation on Tyr 142 by the kinase activity of WSTF (Williams-Beuren syndrome transcription factor), which prevents MDC1 focus formation and instead binds the proapoptotic kinase JNK1 (Cook et al. 2009; Xiao et al. 2009). This regulatory mechanism is thought to govern the balance between DNA damage signaling and cell death.

Once bound to γ H2AX, MDC1 in turn acts as a loading platform for other DDR components, with many of these interactions also being phospho-dependent (Jungmichel and Stucki 2010). For instance, ATM-dependent phosphorylation of MDC1 on Thr-Gln-X-Phe (TQXF) motifs creates binding sites for the FHA domain of the ubiquitin E3 ligase RNF8 (Ring finger protein 8), which in turn promotes the focal accumulation of 53BP1 and BRCA1 at DSB sites (Huen et al. 2007; Kolas et al. 2007; Mailand et al. 2007). Furthermore, constitutive phosphorylation of

Polo and Jackson




Figure 6. Specialized binding modules for recognition of post-translational modifications (PTMs) at DNA breaks. The recruitment of DDR proteins to modified histones or other modified proteins at sites of DNA breaks is mediated by specific interactions between the post-translational modification and a dedicated binding module. BRCT and FHA domains, which are represented by red and green semicircles, bind phosphorylated serine or threonine residues; Tudor domains, chromodomains, and PDH finger domains bind methylated histones; bromodomains (Bromo) bind acetylated histones; and UBDs bind ubiquitylated proteins. The PAR-binding domain can take the form of a basic stretch of amino acids (Basic), a PAR-binding zinc finger (PBZ), or a macrodomain (Macro). Note that some of these modules are found as tandem domains and that not all post-translational modifications are damage-induced (asterisk [*] denotes constitutive modifications). The species of the proteins are indicated, unless only human proteins are listed. (*H.s.*) *Homo sapiens*; (*S.c.*) *S. cerevisiae*, (*S.p.*) *S. pombe*.

MDC1 by CK2 (casein kinase 2) on Ser–Asp–Thr–Asp (SDTD) repeat motifs mediates DSB focus formation by MRN (Fig. 5; Chapman and Jackson 2008; Melander et al. 2008; Spycher et al. 2008; Wu et al. 2008). The molecular basis for this MDC1–NBS1 interaction was unveiled recently by structural and biochemical studies that revealed the existence of a compact and evolutionarily conserved phospho-protein interaction module in NBS1 formed by its closely apposed FHA and BRCT domains (Lloyd et al. 2009; Williams et al. 2009; Hari et al. 2010). Interestingly, while this module mediates binding to MDC1 phospho-SDTD repeats in human cells, it binds the phosphorylated CtlP ortholog in fission yeast and interacts with the phosphorylated XRCC4 ortholog in budding yeast (Matsuzaki et al. 2008; Palmboos et al. 2008; Lloyd et al. 2009; Williams et al. 2009). Some additional FHA- and BRCT-mediated interactions involved in the phospho-dependent recruitment and/or retention of DDR factors at DNA breaks are illustrated in Figure 6.

Notably, in some cases, phosphorylation promotes the dissociation of proteins from sites of DNA breaks. The first characterized example of this was provided by the demonstration that DNA-PKcs autophosphorylation causes it to dissociate from Ku (Chan and Lees-Miller 1996; Merkle et al. 2002). Another example of a phospho-dependent dissociation mechanism is provided by the release from chromatin of the transcriptional cofactor KAP1, which depends on its phosphorylation by ATM (Goodarzi et al. 2008). Similarly, ATM and ATR-mediated phosphorylation of Chk1 is linked to Chk1 dissociation from chromatin in response to DNA damage in mamma-

lian cells, presumably to allow Chk1 to access downstream target proteins in the nucleoplasm (Smits et al. 2006). In fission yeast, hyperphosphorylation of Rad9, which is part of the 9-1-1 checkpoint complex, causes it to dissociate from damaged chromatin by loosening its interaction with RPA (Furuya et al. 2010). This phospho-dependent dissociation mechanism seems critical for efficient repair of DNA damage, and thus potentially contributes to the transition from DNA damage signaling to repair. Furthermore, delocalization of the heterochromatin component HP1 from DNA damage sites has been reported to take place upon CK2-dependent phosphorylation within its chromodomain (Ayoub et al. 2008). This disrupts HP1 interactions with heterochromatin marks, and could thereby contribute to relieving the inhibitory effect of chromatin compaction on the DDR. Notably, however, HP1 recruitment to DNA breaks (and other types of DNA lesions) has also been observed, suggesting more active and dynamic roles for HP1 in the DDR (Ayoub et al. 2008, 2009; Dinant and Luijsterburg 2009; Luijsterburg et al. 2009; Zarebski et al. 2009). Future work will be necessary to fully elucidate HP1 dynamics at sites of DNA breaks, its underlying mechanisms, and its biological functions.

Ubiquitin-dependent signaling mediates DDR focus assembly

Ubiquitylation is the process whereby the 76-amino-acid polypeptide ubiquitin is covalently attached to other proteins singly (monoubiquitylation) or in the form of polyubiquitin chains (polyubiquitylation) by the concerted

actions of ubiquitin E1, E2, and E3 ligase proteins (Pickart 2001). Recent studies have revealed the critical importance of such events in orchestrating the assembly of DDR proteins at DSB sites in vertebrate cells (Messick and Greenberg 2009; Al-Hakim et al. 2010), although whether similar mechanisms also operate in yeast remains to be determined. For example, various protein-ubiquitin conjugates, including ubiquitylated histones, have been detected at sites of DNA breaks in mammalian cells (Morris and Solomon 2004; Polanowska et al. 2006; Mailand et al. 2007; Doil et al. 2009; Stewart et al. 2009). While it is possible that some of these modifications target the associated protein for proteasome-mediated degradation, most do not, and instead serve as docking platforms for focal DDR-protein assembly. Consistent with ubiquitylation actively taking place at such locations, several ubiquitin ligases have been shown to accumulate at sites of DNA breaks in mammalian cells, including BRCA1, RNF8, RNF168, RAD18, HERC2, and PRC1 (Polycomb-repressive complex 1) (Scully et al. 1997; Huen et al. 2007; Kolas et al. 2007; Mailand et al. 2007; Wang and Elledge 2007; Doil et al. 2009; Huang et al. 2009b; Stewart et al. 2009; Watanabe et al. 2009; Wu et al. 2009a; Bekker-Jensen et al. 2010; Chou et al. 2010).

BRCA1 and its interacting partner, BARD1 (BRCA1-associated RING domain), comprise the first mammalian E3 enzyme shown to function within foci at sites of DNA breaks (Hashizume et al. 2001; Morris and Solomon 2004). While few BRCA1 ubiquitylation targets have been identified so far, one of these is CtIP, with BRCA1-mediated ubiquitylation promoting CtIP binding to damaged chromatin (Yu et al. 2006). Besides work on BRCA1/BARD1, much attention has focused on ubiquitylation of histones H2A, H2B, and H2AX, and possibly other substrates by the ubiquitin E3 ligase RNF8, which promotes the focal recruitment of various DDR factors to sites of DNA strand breaks. These factors include the checkpoint mediators 53BP1, BRCA1, and PTIP (Pax2 transactivation domain interaction protein) (Huen et al. 2007; Kolas et al. 2007; Mailand et al. 2007; Wang and Elledge 2007; Doil et al. 2009; Gong et al. 2009; Stewart et al. 2009; Wu et al. 2009a,b); the ubiquitin ligase RAD18, which elicits post-replication repair of damaged DNA (Huang et al. 2009b); and the histone chaperone NPM1 (nucleophosmin) (Koike et al. 2010). In a parallel pathway, H2AX ubiquitylation by PRC1 also contributes to 53BP1 and BRCA1 recruitment to DSBs (Ismail et al. 2010). It will be important to clarify whether and how such PRC1-dependent events cross-talk with the RNF8-dependent pathway. RNF8-dependent ubiquitylation is further amplified by the RNF168 ubiquitin ligase, which is mutated in patients suffering from RIDDLE syndrome (radiosensitivity, immunodeficiency, dysmorphic features, and learning difficulties) (Doil et al. 2009; Pinato et al. 2009; Stewart et al. 2009). Notably, both RNF8 and RNF168 function with the ubiquitin-conjugating E2 enzyme UBC13, and the ubiquitin E3 ligase HERC2 was shown recently to promote RNF8 assembly with UBC13, as opposed to other ubiquitin E2 enzymes (Bekker-Jensen et al. 2010). Additionally, HERC2 stabilizes RNF168 protein levels, thereby regulating ubiquitin-dependent re-

tention of 53BP1 and BRCA1 at DSB sites (Bekker-Jensen et al. 2010).

Analogous to the situation with various DDR-protein phosphorylations, chains of ubiquitin at sites of DNA breaks can be recognized by specific protein domains, termed ubiquitin-binding domains (UBDs) (Fig. 6). While such motifs have been identified in the BRCA1-associated protein RAP80 (receptor-associated protein 80) (H Kim et al. 2007; Sobhian et al. 2007; Sato et al. 2009; Sims and Cohen 2009), the ubiquitin ligase RNF168 (Doil et al. 2009; Pinato et al. 2009, 2011; Stewart et al. 2009), and the RAD18 zinc finger domain (Huang et al. 2009b), the molecular details of ubiquitin-mediated 53BP1 and PTIP recruitment to DNA damage sites remain elusive and probably involve additional post-translational modifications. Indeed, PTIP focus formation is BRCT-dependent (Manke et al. 2003), suggesting a requirement for phosphorylation, while, as described further below, 53BP1 recruitment depends on its ability to bind methylated histone residues (Huyen et al. 2004; Botuyan et al. 2006). In this respect, it is noteworthy that histone ubiquitylation and methylation can act cooperatively, since histone H4 ubiquitylation by the ubiquitin ligase BBAP (B-lymphoma and BAL-associated protein) was shown to stimulate chromatin association of an H4 methylase, thus promoting 53BP1 recruitment to damage sites (Yan et al. 2009). Besides ubiquitylation at DNA damage sites recruiting DDR components to such regions, ubiquitylation of a DDR factor can conversely serve to target this factor to DNA breaks. For instance, when DNA breaks occur during DNA replication, the FA core complex monoubiquitylates the FA proteins FANCD2 and FANCI in a manner that promotes their localization to HR foci (Alpi and Patel 2009; Moldovan and D'Andrea 2009). Ubiquitylated FA proteins in turn recruit the recently identified FAN1 nuclease to sites of DNA repair in a manner dependent on the FAN1 UBD, which might contribute to the resolution of recombination intermediates (Kratz et al. 2010; T Liu et al. 2010; MacKay et al. 2010; Smogorzewska et al. 2010).

Sumoylation controls DDR focus assembly

Modification of proteins by SUMO (small ubiquitin-like modifier) (Creton and Jentsch 2010) has also emerged as an important regulator of the DDR (Bergink and Jentsch 2009), as demonstrated in both yeast and mammalian cells. For example, numerous DDR proteins—including human BLM, XRCC4, and RPA, and budding yeast Sgs1, Ku70, and Rad52—have been identified as sumoylation targets (Eladad et al. 2005; Zhao and Blobel 2005; Branzei et al. 2006; Sacher et al. 2006; Yurchenko et al. 2006; Dou et al. 2010). Furthermore, it was established recently that SUMO conjugation takes place at DNA break sites in mammalian cells, and that the ensuing accumulation of sumoylated proteins in these regions then promotes both protein ubiquitylation at DNA break sites and DDR focus formation (Galanty et al. 2009; Morris et al. 2009). More specifically, these studies revealed that sumoylation of 53BP1 and BRCA1 by the PIAS1 and PIAS4 SUMO E3 ligases increases the residence times of 53BP1 and BRCA1

at damage foci, and serves to enhance BRCA1/BARD1 ubiquitin E3 ligase activity. Other DDR proteins identified as sumoylation targets in response to DNA damage include budding yeast Rad52 (Sacher et al. 2006), whose sumoylation regulates its focal accumulation and recombination activity (Sacher et al. 2006; Torres-Rosell et al. 2007; Altmannova et al. 2010), and the human RPA70 subunit, whose sumoylation enhances its binding to Rad51 and facilitates Rad51 recruitment to DNA damage foci (Dou et al. 2010). Notably, sumoylation is also involved in targeting unrepaired broken DNA ends to the nuclear periphery in yeast (Nagai et al. 2008; Kalocsay et al. 2009), although it is not yet known if sumoylation promotes analogous events in mammalian cells.

Protein methylation-dependent events at DDR foci

Besides requiring histone and nonhistone protein phosphorylation and ubiquitylation, the focal recruitment of 53BP1 and its yeast orthologs to DNA breaks involves their tandem Tudor domains recognizing methylated histone residues (Fig. 6). Specifically, fission yeast Crb2 binds H4K20me2 (Sanders et al. 2004; Greeson et al. 2008) and budding yeast Rad9 binds H3K79me3 (Grenon et al. 2007), while human 53BP1 was initially reported to bind H3K79me3 and subsequently H4K20me1/2 (mono or dimethylated), which might be more physiologically relevant (Huyen et al. 2004; Botuyan et al. 2006; Oda et al. 2010). Importantly, it seems that these histone modifications are constitutive (Huyen et al. 2004; Sanders et al. 2004), which is different from H2AX phosphorylation and histone ubiquitylation that are DNA damage-induced. A proposed model is that these methylated histone residues are not normally readily accessible by 53BP1 and its counterparts, but become exposed upon DNA damage-induced changes in chromatin conformation resulting from histone ubiquitylation (Huen and Chen 2010). Nevertheless, this model is still speculative and might need to be revised, particularly in light of recent reports showing that the H4 methylase PR-Set7/Set8 is recruited to DNA damage sites, and that the H4K20me1/2 it directs is induced to some extent after DNA damage induction in human cells (Yan et al. 2009; Oda et al. 2010).

Tudor domains, PHD, and chromodomains can also recognize methylated lysine residues and target DDR factors to sites of DNA breaks (Fig. 6). For example, H3K4me3 binding at antigen receptor genes by the RAG2 recombinase PHD finger is necessary for effective V(D)J recombination, a programmed gene rearrangement that occurs during B-cell and T-cell development (Liu et al. 2007; Matthews et al. 2007). Furthermore, the chromodomain of the chromatin-modifying complex protein Tip60 binds H3K9me3 when HP1 dissociates from damage sites (Sun et al. 2009). In this case, however, binding to methylated histones acts as an allosteric regulator of Tip60 acetyltransferase function rather than directly promoting Tip60 recruitment to DSBs, which is instead mediated by MRN and ATM. Methylation of DDR factors themselves can also regulate their focal recruitment to DNA breaks. For instance, methylation of MRE11 and 53BP1 on GAR

(glycine–arginine-rich) motifs promotes their focal accumulation by regulating their DNA-binding activities (Boisvert et al. 2005a,b; Dery et al. 2008).

Protein acetylation regulates DDR factor assembly

The importance of a tightly controlled histone acetylation status near DSBs is underlined by the recruitment of several histone acetyltransferases (Gcn5, Esa1, Hat1, and NuA4) and deacetylases (Rpd3, Sir2, and Hst1) to HO-induced DSBs in budding yeast (Downs et al. 2004; Tamburini and Tyler 2005; Qin and Parthun 2006). Similarly, the recruitment of the Tip60 acetyltransferase and the deacetylase enzymes HDAC1, HDAC2, HDAC4, SIRT1, and SIRT6 has been observed at DSB sites in mammalian cells (Kao et al. 2003; Murr et al. 2006; O'Hagan et al. 2008; Oberdoerffer et al. 2008; Kaidi et al. 2010; Miller et al. 2010), and several studies have suggested roles for histone acetylation in regulating the dynamics of DDR factors in the vicinity of DNA breaks. For instance, SIRT1 binding in the vicinity of a DSB promotes recruitment of NBS1 and RAD51 (Oberdoerffer et al. 2008), while SIRT6-dependent deacetylation of the CtIP protein in response to DSBs stimulates RPA and RAD51 focus formation, thus promoting ATR signaling and DSB repair by HR (Kaidi et al. 2010). On the other hand, H3K56 deacetylation by HDAC1 and HDAC2 regulates binding of NHEJ factors to DSB regions (Miller et al. 2010). Additionally, MOF (males absent on the first)-dependent acetylation of H4K16 is important for IR-induced focus formation of MDC1, 53BP1, and BRCA1 in mammalian cells, possibly through interactions between the histone H4 tail and H2AX stimulating the binding of MDC1 to γ H2AX (X Li et al. 2010; Sharma et al. 2010). Along the same lines, H4 acetylation by the MOF-like protein Tip60 at an I-SceI-induced DNA break stimulates the loading of 53BP1 and BRCA1 at such locations (Murr et al. 2006). Tip60 also acetylates H2AX, which promotes H2AX eviction from damaged chromatin, as shown in both *Drosophila* and mammalian cells (Kusch et al. 2004; Ikura et al. 2007). Furthermore, the acetylation status of histone proteins in the vicinity of DNA breaks can regulate the assembly of DDR factors indirectly by modulating chromatin compaction, or directly as exemplified by DNA damage-induced acetylation of histone H3 by Gcn5, which then recruits the chromatin remodeling factor BRG1 by creating a binding site for the BRG1 bromodomain (Fig. 6; Lee et al. 2010).

PARylation targets DDR proteins to DNA breaks

The covalent modification of proteins with ADP-ribose polymers, a process known as PARylation, is catalyzed by PARP enzymes (Hakme et al. 2008). Although PARP enzymes do not exist in the commonly used yeast model systems of *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*, PARP proteins comprise a large family broadly distributed among eukaryotes, several members of which have clearly identified DDR functions (Citarelli et al. 2010). Imposed by PARP1 and PARP2 enzymes in response to DNA breaks in mammalian cells, where it has been most thoroughly studied, PARylation is one of

the earliest events in the DDR but is quickly removed by the action of PARG (PAR glycohydrolase) (Gagne et al. 2006; Hakme et al. 2008; Krishnakumar and Kraus 2010). Through the use of anti-PAR antibodies, PARylation can be detected locally at sites of DNA breaks in mammalian cells (Tartier et al. 2003), where it promotes recruitment of the DNA break repair factors XRCC1 (El-Khamisy et al. 2003; Okano et al. 2003) and APLF (Bekker-Jensen et al. 2007; Kanno et al. 2007; Ahel et al. 2008; Rulten et al. 2008; Eustermann et al. 2010; GY Li et al. 2010). PARylation at DNA breaks is also required for accrual of the chromatin remodeling factors ALC1 and CHD4 (Ahel et al. 2009; Gottschalk et al. 2009; Chou et al. 2010; Polo et al. 2010), the Polycomb histone-modifying complex (Chou et al. 2010), and the histone variant macroH2A (Timinszky et al. 2009). Current models envision these PAR-dependent recruitment events modulating chromatin structure locally at sites of DNA breaks in order to facilitate DNA damage signaling and/or repair. A contribution of PARylation to the early recruitment of MRN has also been reported (Haince et al. 2008). In each of the above cases, specific domains or motifs on DDR factors mediate their binding to PAR (Fig. 6; Pleschke et al. 2000; Karras et al. 2005; Ahel et al. 2008; Gagne et al. 2008; Rulten et al. 2008; Eustermann et al. 2010; Isogai et al. 2010). Interestingly, PARylation can also promote protein dissociation from DNA damage, as shown for the histone chaperone FACT (facilitates chromatin transcription), which is released from damaged chromatin upon PARylation of its Spt16 subunit (Huang et al. 2006; Heo et al. 2008).

Focus disassembly by reversion of post-translational modifications

Resumption of cell cycle progression following DNA repair requires switching off the DDR (Bartek and Lukas 2007), which involves disassembly of DDR foci. In many cases, this appears to occur mainly by reversing the post-translational modifications that led to focal DDR protein assembly. For example, PARG is responsible for erasing PARylation, as described in the preceding section (Gagne et al. 2006). Additionally, several γ H2AX phosphatases have been identified, including Pph3 and PP1 in budding yeast (Keogh et al. 2006; Bazzi et al. 2010) and PP1, PP2A, PP4, PP6, and WIP1 in mammals (Nazarov et al. 2003; Chowdhury et al. 2005, 2008; Nakada et al. 2008; Cha et al. 2010; Douglas et al. 2010; Macurek et al. 2010; Moon et al. 2010). Interestingly, H2AX dephosphorylation, which plays an important role in terminating checkpoint signaling, can take place on chromatin, as reported in human cells (Chowdhury et al. 2005, 2008; Nakada et al. 2008), or after histone eviction from chromatin in yeast (Keogh et al. 2006). Furthermore, the reversal of H2AX phosphorylation also involves Tip60-dependent histone acetylation and subsequent histone eviction from damaged chromatin in *Drosophila* and human cells (Kusch et al. 2004; Jha et al. 2008).

Acting in a manner analogous to phosphatases, deubiquitylating enzymes (DUBs) have also been implicated in

terminating DDR processes. DUBs that function in this regard include USP3 (ubiquitin-specific protease 3), which deubiquitylates histone H2A and H2B (Nicassio et al. 2007), and BRCC36 (BRCA1/BRCA2-containing complex subunit 36), which antagonizes RNF8-dependent ubiquitylation events at DSBs (Shao et al. 2009). Notably, it was shown recently that OTUB1 (OTU domain ubiquitin aldehyde binding 1) also suppresses RNF168-dependent ubiquitylation at DSBs, although in this case this occurs independently of OTUB1 catalytic activity through direct inhibition of the ubiquitin-conjugating enzyme UBC13 (Nakada et al. 2010). In addition, USP16-mediated deubiquitylation of histone H2A was shown to relieve the inhibition of RNA polymerase II transcription at DSBs (Shanbhag et al. 2010), while USP1 counteracts FA protein ubiquitylation (Nijman et al. 2005). Another DUB that regulates the DDR is USP28, which in part operates by stabilizing several DDR factors (Zhang et al. 2006). Although it remains to be demonstrated, it seems likely that desumoylating enzymes will also play important roles in promoting DDR foci disassembly after DNA repair is complete.

Intriguingly, in some cases, DDR factor automodification is coupled to its dissociation from DNA damage sites. For instance, DNA-PKcs autophosphorylation induces a significant conformational change in the protein that elicits its disassembly from Ku and the associated DNA DSB (Chan and Lees-Miller 1996; Merkle et al. 2002; Uematsu et al. 2007; Hammel et al. 2010). Similarly, auto-PARylation of PARP-1 stimulates its dissociation from DNA damage sites, which probably at least partly arise from electrostatic repulsions between DNA and the highly negatively charged PAR chains (Ferro and Olivera 1982; Mortusewicz et al. 2007). Given that acetyltransferases, ubiquitylating enzymes, and sumoylating enzymes are capable of automodifications, it is conceivable that additional examples of automodification-triggered DDR focus disassembly await identification.

Functional importance of DDR foci

The direct binding of DNA breaks by factors such as Ku and MRN is clearly crucial for the DNA repair events that they control. Similarly, the recruitment and activation of the apical DDR kinases ATM, ATR, and DNA-PKcs have well-defined roles at sites of DNA breaks and in DDR focus formation. In contrast, the functional importance of recruiting and activating downstream DDR factors is less well understood and has not been straightforward to decipher. Key reasons for this lack of understanding are the sheer complexity and diversity of downstream DDR responses, and the fact that multiple systems appear to cooperate to control the formation of DDR foci, some of which likely have overlapping and/or compensatory functions. Nevertheless, as summarized below, experimental data and evolutionary conservation indicate that proteins involved in DDR focus assembly and the activities that DDR foci control are of major importance for the maintenance of genome stability.

Evolutionary conservation and biological significance of DDR factor assembly

Many of the proteins and protein modules directing the focal assembly of DDR factors have been highly conserved throughout eukaryotic evolution. A prime example of this is illustrated by the H2AX C-terminal sequence and the tandem BRCT domains that recognize γ H2AX, both of which are conserved from yeasts to humans with only a very few known exceptions (Lee et al. 2005; Stucki et al. 2005; Hammet et al. 2007; Bonner et al. 2008; Kilkenny et al. 2008). Similarly, the Tudor domains of human 53BP1 are also found in yeast 53BP1 orthologs, and associate with similar or identical methylated histone residues in all species examined (Huyen et al. 2004; Sanders et al. 2004; Botuyan et al. 2006; Grenon et al. 2007; Greeson et al. 2008; Oda et al. 2010). Other examples of strong evolutionary conservation are provided by the checkpoint kinases, which occur in virtually all eukaryotes studied and appear to rely on highly conserved mechanisms for their recruitment and activation in response to DNA breaks (Falck et al. 2005; Stracker et al. 2009). Although they have been less studied, it seems likely that the functions of chromatin-modifying proteins, acetyltransferases, deacetylases, PARP enzymes, and components of the ubiquitylation and sumoylation systems in controlling DDR focus assembly will also turn out to be highly conserved, at least among metazoans.

Not only are focus-forming DDR factors and their recruitment mechanisms evolutionarily conserved, but defects in such proteins are associated with various pathologies, in both model systems and human genetic conditions. For instance, key regulators of DDR focus formation—such as ATM, ATR, MRE11, NBS1, and RNF168—are mutated in severe genome instability disorders (Jackson and Bartek 2009; Ciccia and Elledge 2010), and MRN, RNF8, and RNF168 have been identified as prominent targets during viral infection (Carson et al. 2009; Lilley et al. 2010). In some cases, however, the ability of certain DDR factors to compensate for one another has hindered evaluation of their functional requirements. A good example of this is provided by mammalian PARP-1 and PARP-2: Single-knockout mice for PARP-1 or PARP-2 are viable, although they display increased genomic instability and hypersensitivity to DNA-damaging agents, while the double knockout is early embryonic-lethal (Wang et al. 1995; de Murcia et al. 1997; Masutani et al. 1999; Menissier de Murcia et al. 2003). Additionally, functional defects caused by DDR factor dysfunction are often not initially obvious but become apparent only under more detailed examination. For instance, while H2AX-deficient mice are viable and show only partial DDR defects, they are growth-retarded and display increased cancer predisposition in the context of oncogenic mutations (Bassing et al. 2002, 2003; Celeste et al. 2002, 2003a), and H2AX-deficient cells show increased error-prone repair of DSBs (Xie et al. 2004). Moreover, while they are viable and overtly normal, mice lacking H2AX, RNF8, or 53BP1, and, to a lesser extent, MDC1, have class switch recombination defects that

would have profound impacts on animal viability in natural environments (Petersen et al. 2001; Celeste et al. 2002; Ward et al. 2004; Lou et al. 2006; Li et al. 2010; Santos et al. 2010).

Structural and regulatory functions of DDR foci

A prominent feature of many proteins that respond to DNA breaks is their ability to concentrate into microscopically detectable foci. It is widely assumed that the focal accumulation of DDR proteins, by increasing their local concentrations, potentiates interactions between them and with the damaged DNA, thus mounting rapid and effective responses to DNA breaks, while at the same time making accidental DDR induction unlikely. Consistent with this idea, studies in yeast and mammalian systems have demonstrated that colocalization of DDR proteins rather than DNA damage per se is critical for DNA damage signaling (Bonilla et al. 2008; Soutoglou and Misteli 2008). Focus-based responses may also have evolved to allow effective DDR events to take place even under conditions in which the individual DDR components are present at fairly low levels in the cell. In this way, initial DDR events would not necessarily rely on other, potentially slower mechanisms to concentrate DDR components in the nucleus, such as by increasing their levels of expression and/or stability. It is also tempting to speculate that increasing the local concentrations of DDR factors at DNA break sites might be particularly critical in chromatin regions otherwise inhibitory to repair, such as heterochromatin (Noon et al. 2010). Another likely key facet of DDR foci is their ability to promote signal amplification through DDR factors spreading along the chromatin surrounding the break. While the importance of such mechanisms might be missed in many experimental settings, where it is common to use high levels of DNA-damaging agents, amplification of DNA damage signaling is likely to be very important in normal biological contexts. This is particularly evident when one considers that DNA damage checkpoints seem to be able to respond to very small numbers of DSBs, with some experimental data indicating that 10–20 DSBs are enough to elicit G2 arrest in human cells (Deckbar et al. 2007), while very few or even a single unrepaired DSB can be sufficient to trigger p53-dependent G1 arrest in human cells (Huang et al. 1996) or cell death in yeast (Bennett et al. 1993).

Along with concentrating DDR factors and amplifying damage signaling, DDR foci may also fulfill more structural roles by stabilizing broken DNA ends and protecting them from excessive degradation and/or illegitimate repair events (Yin et al. 2009). In line with this idea, structural maintenance of chromosome (SMC) proteins, including cohesins, are recruited to chromosomes in response to DSBs in budding yeast, where they facilitate DSB repair by HR by maintaining sister chromatids in close proximity (Strom et al. 2004; Unal et al. 2004, 2007; De Piccoli et al. 2006; Lindroos et al. 2006). Tethering of DNA ends together can also be achieved through the dimerization or oligomerization of DDR proteins, as shown for the MRN complex (de Jager et al. 2001; Moreno-Herrero et al. 2005) and

53BP1, which promotes long-range joining of deprotected telomeres and V(D)J recombination intermediates in mammalian cells (Difilippantonio et al. 2008; Dimitrova et al. 2008). Aside from such structural functions, DDR foci can also be viewed as a temporary storage site for some DDR factors, as illustrated in mitotic cells, where they mark sites of DNA damage for full DDR activation only in the following G1 phase (Giunta et al. 2010). Another example of a specialized DSB storage focus is the telomere-associated Shelterin complex, which sequesters chromosomal termini and prevents them from being recognized as damaged substrates by forming a protective complex at telomeres after their replication (Verdun et al. 2005). Another likely regulatory function for DDR foci is to contribute to the proper coordination of DNA damage signaling and repair with other DNA metabolic activities by inhibiting replication and transcription. In this regard, DNA methylation and histone modifications—including deacetylation, methylation, and ubiquitylation at sites of DNA breaks—have been proposed to contribute to silencing of damaged chromatin (O'Hagan et al. 2008; Shanbhag et al. 2010).

Conclusions and future directions

The focal accumulation of DDR proteins at sites of DNA breaks was first described >15 years ago, and the molecular mechanisms of focus formation have been the subject of intense investigation since then. As a result, we have identified many of the key protein players in these events and have a growing understanding of their biochemical and physiological functions. In particular, considerable advances have been made in elucidating the hierarchical and functional organization of DDR factor assemblies at DNA break sites, which we now know represent intricate protein and nucleic acid networks precisely coordinated in space and time. Furthermore, recent studies on the dynamics of DDR foci have revealed the prominent roles of combinatorial protein modifications in ensuring rapid, reversible, and fine-tuned regulation of DDR factor recruitment to and dissociation from sites of DNA breaks.

Nevertheless, there are many remaining challenges that will surely attract the attentions of DDR researchers for many years to come. First, while many DDR components are known, the frequent reporting of additional factors in the literature suggests that many more await identification. In this regard, ongoing genetic, proteomic, and siRNA-based screens seem set to provide many additional DDR components and regulators whose functions must then be defined. Another major challenge for the future will be to understand not only how the various post-translational modifications are imposed on DDR factors and which binding modules they specifically associate with, but also how and when they get erased to allow termination of the DDR. It will also be of great interest to further explore likely cross-talks—cooperative or antagonistic—between these post-translational modifications in coordinating the assembly/disassembly of DDR factors at DNA breaks, which we are only beginning to uncover. Further exciting challenges include determining how DNA break signaling and repair are influenced by cell physiology and disease,

and how they may be differentially regulated in different cell types and tissues.

As with many areas of biological research, recent progress in the DDR field has been highly dependent on the development of new methodologies and equipment, including cutting-edge microscopy techniques. Thus, it seems that further progress in the field is also likely to be triggered by further technical improvements, such as new superresolution microscopes and associated computational tools that will allow hitherto unattainable resolution of DDR foci in both three-dimensional space and time (Schermele et al. 2008). Complementing such work will be the development of further biochemical and structural analyses of DDR components, together with siRNA-based and genetic manipulation techniques in cells and model organisms.

Finally, we emphasize that DNA break-associated proteins and the foci that they assemble into are of considerable medical importance, with defects in them being associated with various pathologies, particularly cancer. In this regard, it is notable that the immunodetection of γ H2AX foci, which indirectly measure DSB formation and repair, is showing promise as a sensitive diagnostic tool to detect cancer cells and also monitor cancer progression and assess responses to treatment (Bonner et al. 2008; Mah et al. 2010). Moreover, the existence of many druggable protein targets in DNA break-associated events is providing exciting opportunities for developing new therapeutic agents that, by exploiting differences between normal cells and cancer cells, have the potential to markedly improve cancer management.

Acknowledgments

We apologize to those whose work we inadvertently did not cite or elected not to cite in order to keep this review to a reasonable length. We thank D. Durocher, R. Scully, K. Miller, and K. Dry for critical reading of the manuscript. Research in the Jackson laboratory is supported by grants from Cancer Research UK, the European Union (GENICA HEALTH-F2-2007-201630), the Wellcome Trust, and the BBSRC, and core infrastructure is provided by funding from Cancer Research UK and the Wellcome Trust. S.P.J. receives his salary from the University of Cambridge with a supplement from Cancer Research UK, while S.E.P. was funded by the Human Frontier Science Program Organization, followed by a fellowship from Association pour la Recherche sur le Cancer.

References

- Ahel I, Ahel D, Matsusaka T, Clark AJ, Pines J, Boulton SJ, West SC. 2008. Poly(ADP-ribose)-binding zinc finger motifs in DNA repair/checkpoint proteins. *Nature* **451**: 81–85.
- Ahel D, Horejsi Z, Wiechens N, Polo SE, Garcia-Wilson E, Ahel I, Flynn H, Skehel M, West SC, Jackson SP, et al. 2009. Poly(ADP-ribose)-dependent regulation of DNA repair by the chromatin remodeling enzyme ALC1. *Science* **325**: 1240–1243.
- Al-Hakim A, Escibano-Diaz C, Landry MC, O'Donnell L, Panier S, Szilard RK, Durocher D. 2010. The ubiquitous role of ubiquitin in the DNA damage response. *DNA Repair (Amst)* **9**: 1229–1240.
- Alpi AF, Patel KJ. 2009. Monoubiquitylation in the Fanconi anemia DNA damage response pathway. *DNA Repair (Amst)* **8**: 430–435.

Polo and Jackson

- Altmannova V, Eckert-Boulet N, Americ M, Kolesar P, Chaloupkova R, Damborsky J, Sung P, Zhao X, Lisby M, Krejci L. 2010. Rad52 SUMOylation affects the efficiency of the DNA repair. *Nucleic Acids Res* **38**: 4708–4721.
- Ame JC, Rolli V, Schreiber V, Niedergang C, Apiou F, Decker P, Muller S, Hoger T, Menissier-de Murcia J, de Murcia G. 1999. PARP-2, a novel mammalian DNA damage-dependent poly(ADP-ribose) polymerase. *J Biol Chem* **274**: 17860–17868.
- Amitani I, Liu B, Dombrowski CC, Baskin RJ, Kowalczykowski SC. 2010. Watching individual proteins acting on single molecules of DNA. *Methods Enzymol* **472**: 261–291.
- Ashley T, Plug AW, Xu J, Solari AJ, Reddy G, Golub EI, Ward DC. 1995. Dynamic changes in Rad51 distribution on chromatin during meiosis in male and female vertebrates. *Chromosoma* **104**: 19–28.
- Ayoub N, Jeyasekharan AD, Bernal JA, Venkitaraman AR. 2008. HP1- β mobilization promotes chromatin changes that initiate the DNA damage response. *Nature* **453**: 682–686.
- Ayoub N, Jeyasekharan AD, Venkitaraman AR. 2009. Mobilization and recruitment of HP1: a bimodal response to DNA breakage. *Cell Cycle* **8**: 2945–2950.
- Ball HL, Ehrhardt MR, Mordes DA, Glick GG, Chazin WJ, Cortez D. 2007. Function of a conserved checkpoint recruitment domain in ATRIP proteins. *Mol Cell Biol* **27**: 3367–3377.
- Barlow JH, Lisby M, Rothstein R. 2008. Differential regulation of the cellular response to DNA double-strand breaks in G1. *Mol Cell* **30**: 73–85.
- Bartek J, Lukas J. 2007. DNA damage checkpoints: from initiation to recovery or adaptation. *Curr Opin Cell Biol* **19**: 238–245.
- Bassing CH, Chua KF, Sekiguchi J, Suh H, Whitlow SR, Fleming JC, Monroe BC, Ciccone DN, Yan C, Vlasakova K, et al. 2002. Increased ionizing radiation sensitivity and genomic instability in the absence of histone H2AX. *Proc Natl Acad Sci* **99**: 8173–8178.
- Bassing CH, Suh H, Ferguson DO, Chua KF, Manis J, Eckersdorff M, Gleason M, Bronson R, Lee C, Alt FW. 2003. Histone H2AX: a dosage-dependent suppressor of oncogenic translocations and tumors. *Cell* **114**: 359–370.
- Bazzi M, Mantiero D, Trovesi C, Lucchini G, Longhese MP. 2010. Dephosphorylation of γ H2A by Glc7/protein phosphatase 1 promotes recovery from inhibition of DNA replication. *Mol Cell Biol* **30**: 131–145.
- Bekker-Jensen S, Lukas C, Kitagawa R, Melander F, Kastan MB, Bartek J, Lukas J. 2006. Spatial organization of the mammalian genome surveillance machinery in response to DNA strand breaks. *J Cell Biol* **173**: 195–206.
- Bekker-Jensen S, Fugger K, Danielsen JR, Gromova I, Sehested M, Celis J, Bartek J, Lukas J, Mailand N. 2007. Human Xip1 (C2orf13) is a novel regulator of cellular responses to DNA strand breaks. *J Biol Chem* **282**: 19638–19643.
- Bekker-Jensen S, Rendtlew Danielsen J, Fugger K, Gromova I, Nerstedt A, Lukas C, Bartek J, Lukas J, Mailand N. 2010. HERC2 coordinates ubiquitin-dependent assembly of DNA repair factors on damaged chromosomes. *Nat Cell Biol* **12**: 80–86.
- Benjamin RC, Gill DM. 1980. Poly(ADP-ribose) synthesis in vitro programmed by damaged DNA. A comparison of DNA molecules containing different types of strand breaks. *J Biol Chem* **255**: 10502–10508.
- Bennett CB, Lewis AL, Baldwin KK, Resnick MA. 1993. Lethality induced by a single site-specific double-strand break in a dispensable yeast plasmid. *Proc Natl Acad Sci* **90**: 5613–5617.
- Bennetzen MV, Larsen DH, Bunkenborg J, Bartek J, Lukas J, Andersen JS. 2010. Site-specific phosphorylation dynamics of the nuclear proteome during the DNA damage response. *Mol Cell Proteomics* **9**: 1314–1323.
- Bensimon A, Schmidt A, Ziv Y, Elkon R, Wang S-Y, Chen DJ, Aebersold R, Shiloh Y. 2010. ATM-dependent and -independent dynamics of the nuclear phosphoproteome after DNA damage. *Sci Signal* **3**: rs3. doi: 10.1126/scisignal.20011034.
- Bergink S, Jentsch S. 2009. Principles of ubiquitin and SUMO modifications in DNA repair. *Nature* **458**: 461–467.
- Berkovich E, Monnat RJ Jr, Kastan MB. 2008. Assessment of protein dynamics and DNA repair following generation of DNA double-strand breaks at defined genomic sites. *Nat Protoc* **3**: 915–922.
- Bernstein E, Hake SB. 2006. The nucleosome: a little variation goes a long way. *Biochem Cell Biol* **84**: 505–517.
- Bernstein KA, Rothstein R. 2009. At loose ends: resecting a double-strand break. *Cell* **137**: 807–810.
- Bernstein KA, Gangloff S, Rothstein R. 2010. The RecQ DNA helicases in DNA repair. *Annu Rev Genet* **44**: 393–417.
- Boisvert FM, Hendzel MJ, Masson JY, Richard S. 2005a. Methylation of MRE11 regulates its nuclear compartmentalization. *Cell Cycle* **4**: 981–989.
- Boisvert FM, Rhie A, Richard S, Doherty AJ. 2005b. The GAR motif of 53BP1 is arginine methylated by PRMT1 and is necessary for 53BP1 DNA binding activity. *Cell Cycle* **4**: 1834–1841.
- Bonilla CY, Melo JA, Toczyski DP. 2008. Colocalization of sensors is sufficient to activate the DNA damage checkpoint in the absence of damage. *Mol Cell* **30**: 267–276.
- Bonner WM, Redon CE, Dickey JS, Nakamura AJ, Sedelnikova OA, Solier S, Pommier Y. 2008. γ H2AX and cancer. *Nat Rev Cancer* **8**: 957–967.
- Bork P, Hofmann K, Bucher P, Neuwald AF, Altschul SF, Koonin EV. 1997. A superfamily of conserved domains in DNA damage-responsive cell cycle checkpoint proteins. *FASEB J* **11**: 68–76.
- Botchway SW, Reynolds P, Parker AW, O'Neill P. 2010. Use of near infrared femtosecond lasers as sub-micron radiation microbeam for cell DNA damage and repair studies. *Mutat Res* **704**: 38–44.
- Botuyan MV, Lee J, Ward IM, Kim JE, Thompson JR, Chen J, Mer G. 2006. Structural basis for the methylation state-specific recognition of histone H4-K20 by 53BP1 and Crb2 in DNA repair. *Cell* **127**: 1361–1373.
- Branzei D, Sollier J, Liberi G, Zhao X, Maeda D, Seki M, Enomoto T, Ohta K, Foiani M. 2006. Ubc9- and mms21-mediated sumoylation counteracts recombinogenic events at damaged replication forks. *Cell* **127**: 509–522.
- Burma S, Chen BP, Murphy M, Kurimasa A, Chen DJ. 2001. ATM phosphorylates histone H2AX in response to DNA double-strand breaks. *J Biol Chem* **276**: 42462–42467.
- Caldecott KW. 2008. Single-strand break repair and genetic disease. *Nat Rev Genet* **9**: 619–631.
- Callebaut I, Mornon JP. 1997. From BRCA1 to RAP1: a widespread BRCT module closely associated with DNA repair. *FEBS Lett* **400**: 25–30.
- Calsou P, Delteil C, Frit P, Drouet J, Salles B. 2003. Coordinated assembly of Ku and p46 subunits of the DNA-dependent protein kinase on DNA ends is necessary for XRCC4-ligase IV recruitment. *J Mol Biol* **326**: 93–103.
- Carson CT, Orazio NI, Lee DV, Suh J, Bekker-Jensen S, Araujo FD, Lakdawala SS, Lilley CE, Bartek J, Lukas J, et al. 2009. Mislocalization of the MRN complex prevents ATR signaling during adenovirus infection. *EMBO J* **28**: 652–662.
- Celeste A, Petersen S, Romanienko PJ, Fernandez-Capetillo O, Chen HT, Sedelnikova OA, Reina-San-Martin B, Coppola V, Meffre E, Difilippantonio MJ, et al. 2002. Genomic instability in mice lacking histone H2AX. *Science* **296**: 922–927.

- Celeste A, Difilippantonio S, Difilippantonio MJ, Fernandez-Capetillo O, Pilch DR, Sedelnikova OA, Eckhaus M, Ried T, Bonner WM, Nussenzweig A. 2003a. H2AX haploinsufficiency modifies genomic stability and tumor susceptibility. *Cell* **114**: 371–383.
- Celeste A, Fernandez-Capetillo O, Kruhlak MJ, Pilch DR, Staudt DW, Lee A, Bonner RF, Bonner WM, Nussenzweig A. 2003b. Histone H2AX phosphorylation is dispensable for the initial recognition of DNA breaks. *Nat Cell Biol* **5**: 675–679.
- Cha H, Lowe JM, Li H, Lee JS, Belova GI, Bulavin DV, Fornace AJ Jr. 2010. Wip1 directly dephosphorylates γ -H2AX and attenuates the DNA damage response. *Cancer Res* **70**: 4112–4122.
- Chadwick BP, Lane TF. 2005. BRCA1 associates with the inactive X chromosome in late S-phase, coupled with transient H2AX phosphorylation. *Chromosoma* **114**: 432–439.
- Chai B, Huang J, Cairns BR, Laurent BC. 2005. Distinct roles for the RSC and Swi/Snf ATP-dependent chromatin remodelers in DNA double-strand break repair. *Genes Dev* **19**: 1656–1661.
- Chan DW, Lees-Miller SP. 1996. The DNA-dependent protein kinase is inactivated by autophosphorylation of the catalytic subunit. *J Biol Chem* **271**: 8936–8941.
- Chapman JR, Jackson SP. 2008. Phospho-dependent interactions between NBS1 and MDC1 mediate chromatin retention of the MRN complex at sites of DNA damage. *EMBO Rep* **9**: 795–801.
- Chen CC, Carson JJ, Feser J, Tamburini B, Zabaronic S, Linger J, Tyler JK. 2008. Acetylated lysine 56 on histone H3 drives chromatin assembly after repair and signals for the completion of repair. *Cell* **134**: 231–243.
- Chou DM, Adamson B, Dephoure NE, Tan X, Nottke AC, Hurov KE, Gygi SP, Colaiacovo MP, Elledge SJ. 2010. A chromatin localization screen reveals poly (ADP ribose)-regulated recruitment of the repressive polycomb and NuRD complexes to sites of DNA damage. *Proc Natl Acad Sci* **107**: 18475–18480.
- Chowdhury D, Keogh MC, Ishii H, Peterson CL, Buratowski S, Lieberman J. 2005. γ -H2AX dephosphorylation by protein phosphatase 2A facilitates DNA double-strand break repair. *Mol Cell* **20**: 801–809.
- Chowdhury D, Xu X, Zhong X, Ahmed F, Zhong J, Liao J, Dykxhoorn DM, Weinstock DM, Pfeifer GP, Lieberman J. 2008. A PP4-phosphatase complex dephosphorylates γ -H2AX generated during DNA replication. *Mol Cell* **31**: 33–46.
- Ciccia A, Elledge SJ. 2010. The DNA damage response: making it safe to play with knives. *Mol Cell* **40**: 179–204.
- Citarelli M, Teotia S, Lamb RS. 2010. Evolutionary history of the poly(ADP-ribose) polymerase gene family in eukaryotes. *BMC Evol Biol* **10**: 308. doi: 10.1186/1471-2148-10-308.
- Clapier CR, Cairns BR. 2009. The biology of chromatin remodeling complexes. *Annu Rev Biochem* **78**: 273–304.
- Clerici M, Mantiero D, Guerini I, Lucchini G, Longhese MP. 2008. The Yku70–Yku80 complex contributes to regulate double-strand break processing and checkpoint activation during the cell cycle. *EMBO Rep* **9**: 810–818.
- Cook PJ, Ju BG, Telese F, Wang X, Glass CK, Rosenfeld MG. 2009. Tyrosine dephosphorylation of H2AX modulates apoptosis and survival decisions. *Nature* **458**: 591–596.
- Cortez D, Guntuku S, Qin J, Elledge SJ. 2001. ATR and ATRIP: partners in checkpoint signaling. *Science* **294**: 1713–1716.
- Cowell IG, Sunter NJ, Singh PB, Austin CA, Durkacz BW, Tilby MJ. 2007. γ H2AX foci form preferentially in euchromatin after ionising-radiation. *PLoS ONE* **2**: e1057. doi: 10.1371/journal.pone.001057.
- Creton S, Jentsch S. 2010. SnapShot: The SUMO system. *Cell* **143**: 848.
- Cuozzo C, Porcellini A, Angrisano T, Morano A, Lee B, Di Pardo A, Messina S, Iuliano R, Fusco A, Santillo MR, et al. 2007. DNA damage, homology-directed repair, and DNA methylation. *PLoS Genet* **3**: e110. doi: 10.1371/journal.pgen.0030110.
- d'Adda di Fagagna F, Reaper PM, Clay-Farrace L, Fiegler H, Carr P, Von Zglinicki T, Saretzki G, Carter NP, Jackson SP. 2003. A DNA damage checkpoint response in telomere-initiated senescence. *Nature* **426**: 194–198.
- D'Amours D, Desnoyers S, D'Silva I, Poirier GG. 1999. Poly(ADP-ribosylation) reactions in the regulation of nuclear functions. *Biochem J* **342**: 249–268.
- de Jager M, van Noort J, van Gent DC, Dekker C, Kanaar R, Wyman C. 2001. Human Rad50/Mre11 is a flexible complex that can tether DNA ends. *Mol Cell* **8**: 1129–1135.
- de Murcia G, Menissier de Murcia J. 1994. Poly(ADP-ribose) polymerase: a molecular nick-sensor. *Trends Biochem Sci* **19**: 172–176.
- de Murcia JM, Niedergang C, Trucco C, Ricoul M, Dutrillaux B, Mark M, Oliver FJ, Masson M, Dierich A, LeMour M, et al. 1997. Requirement of poly(ADP-ribose) polymerase in recovery from DNA damage in mice and in cells. *Proc Natl Acad Sci* **94**: 7303–7307.
- Deckbar D, Birraux J, Krempler A, Tchouandong L, Beucher A, Walker S, Stiff T, Jeggo P, Lobrich M. 2007. Chromosome breakage after G2 checkpoint release. *J Cell Biol* **176**: 749–755.
- De Piccoli G, Cortes-Ledesma F, Ira G, Torres-Rosell J, Uhle S, Farmer S, Hwang JY, Machin F, Ceschia A, McAleenan A, et al. 2006. Smc5–Smc6 mediate DNA double-strand-break repair by promoting sister-chromatid recombination. *Nat Cell Biol* **8**: 1032–1034.
- Dery U, Coulombe Y, Rodrigue A, Stasiak A, Richard S, Masson JY. 2008. A glycine-arginine domain in control of the human MRE11 DNA repair protein. *Mol Cell Biol* **28**: 3058–3069.
- Difilippantonio S, Gapud E, Wong N, Huang CY, Mahowald G, Chen HT, Kruhlak MJ, Callen E, Livak F, Nussenzweig MC, et al. 2008. 53BP1 facilitates long-range DNA end-joining during V(D)J recombination. *Nature* **456**: 529–533.
- Dimitrova N, Chen YC, Spector DL, de Lange T. 2008. 53BP1 promotes non-homologous end joining of telomeres by increasing chromatin mobility. *Nature* **456**: 524–528.
- Dinant C, Luijsterburg MS. 2009. The emerging role of HP1 in the DNA damage response. *Mol Cell Biol* **29**: 6335–6340.
- Doil C, Mailand N, Bekker-Jensen S, Menard P, Larsen DH, Pepperkok R, Ellenberg J, Panier S, Durocher D, Bartek J, et al. 2009. RNF168 binds and amplifies ubiquitin conjugates on damaged chromosomes to allow accumulation of repair proteins. *Cell* **136**: 435–446.
- Dou H, Huang C, Singh M, Carpenter PB, Yeh ET. 2010. Regulation of DNA repair through deSUMOylation and SUMOylation of replication protein A complex. *Mol Cell* **39**: 333–345.
- Douglas P, Zhong J, Ye R, Moorhead GB, Xu X, Lees-Miller SP. 2010. Protein phosphatase 6 interacts with the DNA-dependent protein kinase catalytic subunit and dephosphorylates γ -H2AX. *Mol Cell Biol* **30**: 1368–1381.
- Downs JA, Lowndes NF, Jackson SP. 2000. A role for *Saccharomyces cerevisiae* histone H2A in DNA repair. *Nature* **408**: 1001–1004.
- Downs JA, Allard S, Jobin-Robitaille O, Javaheri A, Auger A, Bouchard N, Kron SJ, Jackson SP, Cote J. 2004. Binding of chromatin-modifying activities to phosphorylated histone H2A at DNA damage sites. *Mol Cell* **16**: 979–990.
- Drouot J, Delteil C, Lefrançois J, Concannon P, Salles B, Calsou P. 2005. DNA-dependent protein kinase and XRCC4-DNA ligase IV mobilization in the cell in response to DNA double strand breaks. *J Biol Chem* **280**: 7060–7069.

Polo and Jackson

- Durocher D, Henckel J, Fersht AR, Jackson SP. 1999. The FHA domain is a modular phosphopeptide recognition motif. *Mol Cell* **4**: 387–394.
- Dvir A, Peterson SR, Knuth MW, Lu H, Dynan WS. 1992. Ku autoantigen is the regulatory component of a template-associated protein kinase that phosphorylates RNA polymerase II. *Proc Natl Acad Sci* **89**: 11920–11924.
- Dynan WS, Yoo S. 1998. Interaction of Ku protein and DNA-dependent protein kinase catalytic subunit with nucleic acids. *Nucleic Acids Res* **26**: 1551–1559.
- Eladad S, Ye TZ, Hu P, Leversha M, Beresten S, Matunis MJ, Ellis NA. 2005. Intra-nuclear trafficking of the BLM helicase to DNA damage-induced foci is regulated by SUMO modification. *Hum Mol Genet* **14**: 1351–1365.
- El-Khamisy SF, Masutani M, Suzuki H, Caldecott KW. 2003. A requirement for PARP-1 for the assembly or stability of XRCC1 nuclear foci at sites of oxidative DNA damage. *Nucleic Acids Res* **31**: 5526–5533.
- Erdel F, Schubert T, Marth C, Langst G, Rippe K. 2010. Human ISWI chromatin-remodeling complexes sample nucleosomes via transient binding reactions and become immobilized at active sites. *Proc Natl Acad Sci* **107**: 19873–19878.
- Essers J, Houtsmuller AB, Kanaar R. 2006. Analysis of DNA recombination and repair proteins in living cells by photo-bleaching microscopy. *Methods Enzymol* **408**: 463–485.
- Eustermann S, Brockmann C, Mehrotra PV, Yang JC, Loakes D, West SC, Ahel I, Neuhaus D. 2010. Solution structures of the two PBZ domains from human APLF and their interaction with poly(ADP-ribose). *Nat Struct Mol Biol* **17**: 241–243.
- Falck J, Coates J, Jackson SP. 2005. Conserved modes of recruitment of ATM, ATR and DNA-PKcs to sites of DNA damage. *Nature* **434**: 605–611.
- Fanti L, Pimpinelli S. 2008. HP1: a functionally multifaceted protein. *Curr Opin Genet Dev* **18**: 169–174.
- Ferro AM, Olivera BM. 1982. Poly(ADP-ribosylation) in vitro. Reaction parameters and enzyme mechanism. *J Biol Chem* **257**: 7808–7813.
- FitzGerald JE, Grenon M, Lowndes NF. 2009. 53BP1: function and mechanisms of focal recruitment. *Biochem Soc Trans* **37**: 897–904.
- Flynn RL, Zou L. 2010. Oligonucleotide/oligosaccharide-binding fold proteins: a growing family of genome guardians. *Crit Rev Biochem Mol Biol* **45**: 266–275.
- Friedberg EC, Walker GC, Siede W, Wood RD, Schultz RA, Ellenberger T. 2006. *DNA repair and mutagenesis*, 2nd ed. ASM Press, New York.
- Furuya K, Miyabe I, Tsutsui Y, Paderi F, Kakusho N, Masai H, Niki H, Carr AM. 2010. DDK phosphorylates checkpoint clamp component Rad9 and promotes its release from damaged chromatin. *Mol Cell* **40**: 606–618.
- Gagne JP, Hendzel MJ, Droit A, Poirier GG. 2006. The expanding role of poly(ADP-ribose) metabolism: current challenges and new perspectives. *Curr Opin Cell Biol* **18**: 145–151.
- Gagne JP, Isabelle M, Lo KS, Bourassa S, Hendzel MJ, Dawson VL, Dawson TM, Poirier GG. 2008. Proteome-wide identification of poly(ADP-ribose) binding proteins and poly(ADP-ribose)-associated protein complexes. *Nucleic Acids Res* **36**: 6959–6976.
- Galanty Y, Belotserkovskaya R, Coates J, Polo S, Miller KM, Jackson SP. 2009. Mammalian SUMO E3-ligases PIAS1 and PIAS4 promote responses to DNA double-strand breaks. *Nature* **462**: 935–939.
- Gell D, Jackson SP. 1999. Mapping of protein–protein interactions within the DNA-dependent protein kinase complex. *Nucleic Acids Res* **27**: 3494–3502.
- Giunta S, Belotserkovskaya R, Jackson SP. 2010. DNA damage signaling in response to double-strand breaks during mitosis. *J Cell Biol* **190**: 197–207.
- Gong Z, Cho YW, Kim JE, Ge K, Chen J. 2009. Accumulation of Pax2 transactivation domain interaction protein (PTIP) at sites of DNA breaks via RNF8-dependent pathway is required for cell survival after DNA damage. *J Biol Chem* **284**: 7284–7293.
- Goodarzi AA, Noon AT, Deckbar D, Ziv Y, Shiloh Y, Lohrlich M, Jeggo PA. 2008. ATM signaling facilitates repair of DNA double-strand breaks associated with heterochromatin. *Mol Cell* **31**: 167–177.
- Gottlieb TM, Jackson SP. 1993. The DNA-dependent protein kinase: requirement for DNA ends and association with Ku antigen. *Cell* **72**: 131–142.
- Gottschalk AJ, Timinszky G, Kong SE, Jin J, Cai Y, Swanson SK, Washburn MP, Florens L, Ladurner AG, Conaway JW, et al. 2009. Poly(ADP-ribosylation) directs recruitment and activation of an ATP-dependent chromatin remodeler. *Proc Natl Acad Sci* **106**: 13770–13774.
- Greeson NT, Sengupta R, Arida AR, Jenuwein T, Sanders SL. 2008. Di-methyl H4 lysine 20 targets the checkpoint protein Crb2 to sites of DNA damage. *J Biol Chem* **283**: 33168–33174.
- Grenon M, Costelloe T, Jimeno S, O’Shaughnessy A, Fitzgerald J, Zgheib O, Degerth L, Lowndes NF. 2007. Docking onto chromatin via the *Saccharomyces cerevisiae* Rad9 Tudor domain. *Yeast* **24**: 105–119.
- Haaf T, Golub EI, Reddy G, Radding CM, Ward DC. 1995. Nuclear foci of mammalian Rad51 recombination protein in somatic cells after DNA damage and its localization in synaptonemal complexes. *Proc Natl Acad Sci* **92**: 2298–2302.
- Haince JF, McDonald D, Rodrigue A, Dery U, Masson JY, Hendzel MJ, Poirier GG. 2008. PARP1-dependent kinetics of recruitment of MRE11 and NBS1 proteins to multiple DNA damage sites. *J Biol Chem* **283**: 1197–1208.
- Hakme A, Wong HK, Dantzer F, Schreiber V. 2008. The expanding field of poly(ADP-ribosylation) reactions. ‘Protein modifications: beyond the usual suspects’ review series. *EMBO Rep* **9**: 1094–1100.
- Hammel M, Yu Y, Mahaney BL, Cai B, Ye R, Phipps BM, Rambo RP, Hura GL, Pelikan M, So S, et al. 2010. Ku and DNA-dependent protein kinase dynamic conformations and assembly regulate DNA binding and the initial non-homologous end joining complex. *J Biol Chem* **285**: 1414–1423.
- Hammet A, Magill C, Heierhorst J, Jackson SP. 2007. Rad9 BRCT domain interaction with phosphorylated H2AX regulates the G1 checkpoint in budding yeast. *EMBO Rep* **8**: 851–857.
- Hari FJ, Spycher C, Jungmichel S, Pavic L, Stucki M. 2010. A divalent FHA/BRCT-binding mechanism couples the MRE11–RAD50–NBS1 complex to damaged chromatin. *EMBO Rep* **11**: 387–392.
- Harper JW, Elledge SJ. 2007. The DNA damage response: ten years after. *Mol Cell* **28**: 739–745.
- Harrison JC, Haber JE. 2006. Surviving the breakup: the DNA damage checkpoint. *Annu Rev Genet* **40**: 209–235.
- Hartlerode AJ, Scully R. 2009. Mechanisms of double-strand break repair in somatic mammalian cells. *Biochem J* **423**: 157–168.
- Hashizume R, Fukuda M, Maeda I, Nishikawa H, Oyake D, Yabuki Y, Ogata H, Ohta T. 2001. The RING heterodimer BRCA1–BARD1 is a ubiquitin ligase inactivated by a breast cancer-derived mutation. *J Biol Chem* **276**: 14537–14540.
- Heo K, Kim H, Choi SH, Choi J, Kim K, Gu J, Lieber MR, Yang AS, An W. 2008. FACT-mediated exchange of histone variant

- H2AX regulated by phosphorylation of H2AX and ADP-ribosylation of Spt16. *Mol Cell* **30**: 86–97.
- Hoeijmakers JH. 2001. Genome maintenance mechanisms for preventing cancer. *Nature* **411**: 366–374.
- Huang LC, Clarkin KC, Wahl GM. 1996. Sensitivity and selectivity of the DNA damage sensor responsible for activating p53-dependent G1 arrest. *Proc Natl Acad Sci* **93**: 4827–4832.
- Huang JY, Chen WH, Chang YL, Wang HT, Chuang WT, Lee SC. 2006. Modulation of nucleosome-binding activity of FACT by poly[ADP-ribosyl]ation. *Nucleic Acids Res* **34**: 2398–2407.
- Huang J, Gong Z, Ghosal G, Chen J. 2009a. SOSS complexes participate in the maintenance of genomic stability. *Mol Cell* **35**: 384–393.
- Huang J, Huen MS, Kim H, Leung CC, Glover JN, Yu X, Chen J. 2009b. RAD18 transmits DNA damage signalling to elicit homologous recombination repair. *Nat Cell Biol* **11**: 592–603.
- Huen MS, Chen J. 2010. Assembly of checkpoint and repair machineries at DNA damage sites. *Trends Biochem Sci* **35**: 101–108.
- Huen MS, Grant R, Manke I, Minn K, Yu X, Yaffe MB, Chen J. 2007. RNF8 transduces the DNA-damage signal via histone ubiquitylation and checkpoint protein assembly. *Cell* **131**: 901–914.
- Huen MS, Sy SM, Chen J. 2010. BRCA1 and its toolbox for the maintenance of genome integrity. *Nat Rev Mol Cell Biol* **11**: 138–148.
- Huertas P. 2010. DNA resection in eukaryotes: deciding how to fix the break. *Nat Struct Mol Biol* **17**: 11–16.
- Huertas P, Jackson SP. 2009. Human CtIP mediates cell cycle control of DNA end resection and double strand break repair. *J Biol Chem* **284**: 9558–9565.
- Huertas P, Cortes-Ledesma F, Sartori AA, Aguilera A, Jackson SP. 2008. CDK targets Sae2 to control DNA-end resection and homologous recombination. *Nature* **455**: 689–692.
- Huyen Y, Zgheib O, Ditullio RA Jr, Gorgoulis VG, Zacharatos P, Petty TJ, Sheston EA, Mellert HS, Stavridi ES, Halazonetis TD. 2004. Methylated lysine 79 of histone H3 targets 53BP1 to DNA double-strand breaks. *Nature* **432**: 406–411.
- Iacovoni JS, Caron P, Lassadi I, Nicolas E, Massip L, Trouche D, Legube G. 2010. High-resolution profiling of γ H2AX around DNA double strand breaks in the mammalian genome. *EMBO J* **29**: 1446–1457.
- Ikura T, Tashiro S, Kakino A, Shima H, Jacob N, Amunugama R, Yoder K, Izumi S, Kuraoka I, Tanaka K, et al. 2007. DNA damage-dependent acetylation and ubiquitination of H2AX enhances chromatin dynamics. *Mol Cell Biol* **27**: 7028–7040.
- Ismail IH, Andrin C, McDonald D, Hendzel MJ. 2010. BMI1-mediated histone ubiquitylation promotes DNA double-strand break repair. *J Cell Biol* **191**: 45–60.
- Isogai S, Kanno S, Ariyoshi M, Tochio H, Ito Y, Yasui A, Shirakawa M. 2010. Solution structure of a zinc-finger domain that binds to poly-ADP-ribose. *Genes Cells* **15**: 101–110.
- Jackson SP, Bartek J. 2009. The DNA-damage response in human biology and disease. *Nature* **461**: 1071–1078.
- Jakob B, Rudolph JH, Gueven N, Lavin MF, Taucher-Scholz G. 2005. Live cell imaging of heavy-ion-induced radiation responses by beamline microscopy. *Radiat Res* **163**: 681–690.
- Jakob B, Splinter J, Taucher-Scholz G. 2009. Positional stability of damaged chromatin domains along radiation tracks in mammalian cells. *Radiat Res* **171**: 405–418.
- Janin M. 1996. Genetic manipulation of genomes with rare-cutting endonucleases. *Trends Genet* **12**: 224–228.
- Jazayeri A, Falck J, Lukas C, Bartek J, Smith GC, Lukas J, Jackson SP. 2006. ATM- and cell cycle-dependent regulation of ATR in response to DNA double-strand breaks. *Nat Cell Biol* **8**: 37–45.
- Jensen RB, Carreira A, Kowalczykowski SC. 2010. Purified human BRCA2 stimulates RAD51-mediated recombination. *Nature* **467**: 678–683.
- Jha S, Shibata E, Dutta A. 2008. Human Rvb1/Tip49 is required for the histone acetyltransferase activity of Tip60/NuA4 and for the downregulation of phosphorylation on H2AX after DNA damage. *Mol Cell Biol* **28**: 2690–2700.
- Jungmichel S, Stucki M. 2010. MDC1: The art of keeping things in focus. *Chromosoma* **119**: 337–349.
- Kaidi A, Weinert BT, Choudhary C, Jackson SP. 2010. Human SIRT6 promotes DNA end resection through CtIP deacetylation. *Science* **329**: 1348–1353.
- Kalocsay M, Hiller NJ, Jentsch S. 2009. Chromosome-wide Rad51 spreading and SUMO-H2A.Z-dependent chromosome fixation in response to a persistent DNA double-strand break. *Mol Cell* **33**: 335–343.
- Kanno S, Kuzuoka H, Sasao S, Hong Z, Lan L, Nakajima S, Yasui A. 2007. A novel human AP endonuclease with conserved zinc-finger-like motifs involved in DNA strand break responses. *EMBO J* **26**: 2094–2103.
- Kao GD, McKenna WG, Guenther MG, Muschel RJ, Lazar MA, Yen TJ. 2003. Histone deacetylase 4 interacts with 53BP1 to mediate the DNA damage response. *J Cell Biol* **160**: 1017–1027.
- Karras GI, Kustatscher G, Buhecha HR, Allen MD, Pugieux C, Sait F, Bycroft M, Ladurner AG. 2005. The macro domain is an ADP-ribose binding module. *EMBO J* **24**: 1911–1920.
- Kashiwaba S, Kitahashi K, Watanabe T, Onoda F, Ohtsu M, Murakami Y. 2010. The mammalian INO80 complex is recruited to DNA damage sites in an ARP8 dependent manner. *Biochem Biophys Res Commun* **402**: 619–625.
- Kent NA, Chambers AL, Downs JA. 2007. Dual chromatin remodeling roles for RSC during DNA double strand break induction and repair at the yeast MAT locus. *J Biol Chem* **282**: 27693–27701.
- Keogh MC, Kim JA, Downey M, Fillingham J, Chowdhury D, Harrison JC, Onishi M, Datta N, Galicia S, Emili A, et al. 2006. A phosphatase complex that dephosphorylates γ H2AX regulates DNA damage checkpoint recovery. *Nature* **439**: 497–501.
- Kilkenny ML, Dore AS, Roe SM, Nestoras K, Ho JC, Watts FZ, Pearl LH. 2008. Structural and functional analysis of the Crb2-BRCT2 domain reveals distinct roles in checkpoint signaling and DNA damage repair. *Genes Dev* **22**: 2034–2047.
- Kim JS, Krasieva TB, Kurumizaka H, Chen DJ, Taylor AM, Yokomori K. 2005. Independent and sequential recruitment of NHEJ and HR factors to DNA damage sites in mammalian cells. *J Cell Biol* **170**: 341–347.
- Kim H, Chen J, Yu X. 2007. Ubiquitin-binding protein RAP80 mediates BRCA1-dependent DNA damage response. *Science* **316**: 1202–1205.
- Kim JA, Kruhlak M, Dotiwala F, Nussenzweig A, Haber JE. 2007. Heterochromatin is refractory to γ -H2AX modification in yeast and mammals. *J Cell Biol* **178**: 209–218.
- Koike A, Nishikawa H, Wu W, Okada Y, Venkitaraman AR, Ohta T. 2010. Recruitment of phosphorylated NPM1 to sites of DNA damage through RNF8-dependent ubiquitin conjugates. *Cancer Res* **70**: 6746–6756.
- Kolas NK, Chapman JR, Nakada S, Ylanko J, Chahwan R, Sweeney FD, Panier S, Mendez M, Wildenhain J, Thomson TM, et al. 2007. Orchestration of the DNA-damage response by the RNF8 ubiquitin ligase. *Science* **318**: 1637–1640.
- Kornberg RD. 1977. Structure of chromatin. *Annu Rev Biochem* **46**: 931–954.

Polo and Jackson

- Kouzarides T. 2007. Chromatin modifications and their function. *Cell* **128**: 693–705.
- Kratz K, Schopf B, Kaden S, Sendoel A, Eberhard R, Lademann C, Cannavo E, Sartori AA, Hengartner MO, Jiricny J. 2010. Deficiency of FANCD2-associated nuclease KIAA1018/FAN1 sensitizes cells to interstrand crosslinking agents. *Cell* **142**: 77–88.
- Krishnakumar R, Kraus WL. 2010. The PARP side of the nucleus: molecular actions, physiological outcomes, and clinical targets. *Mol Cell* **39**: 8–24.
- Kruhlak MJ, Celeste A, Dellaire G, Fernandez-Capetillo O, Muller WG, McNally JG, Bazett-Jones DP, Nussenzweig A. 2006. Changes in chromatin structure and mobility in living cells at sites of DNA double-strand breaks. *J Cell Biol* **172**: 823–834.
- Kruhlak M, Crouch EE, Orlov M, Montano C, Gorski SA, Nussenzweig A, Misteli T, Phair RD, Casellas R. 2007. The ATM repair pathway inhibits RNA polymerase I transcription in response to chromosome breaks. *Nature* **447**: 730–734.
- Kulis M, Esteller M. 2010. DNA methylation and cancer. *Adv Genet* **70**: 27–56.
- Kusch T, Florens L, Macdonald WH, Swanson SK, Glaser RL, Yates JR III, Abmayr SM, Washburn MP, Workman JL. 2004. Acetylation by Tip60 is required for selective histone variant exchange at DNA lesions. *Science* **306**: 2084–2087.
- Larsen DH, Poinssignon C, Gudjonsson T, Dinant C, Payne MR, Hari FJ, Danielsen JM, Menard P, Sand JC, Stucki M, et al. 2010. The chromatin-remodeling factor CHD4 coordinates signaling and repair after DNA damage. *J Cell Biol* **190**: 731–740.
- Lavin MF. 2007. ATM and the Mre11 complex combine to recognize and signal DNA double-strand breaks. *Oncogene* **26**: 7749–7758.
- Lazzaro F, Giannattasio M, Puddu F, Granata M, Pelliccioli A, Plevani P, Muzi-Falconi M. 2009. Checkpoint mechanisms at the intersection between DNA damage and repair. *DNA Repair (Amst)* **8**: 1055–1067.
- Lee JH, Paull TT. 2005. ATM activation by DNA double-strand breaks through the Mre11–Rad50–Nbs1 complex. *Science* **308**: 551–554.
- Lee MS, Edwards RA, Thede GL, Glover JN. 2005. Structure of the BRCT repeat domain of MDC1 and its specificity for the free COOH-terminal end of the γ -H2AX histone tail. *J Biol Chem* **280**: 32053–32056.
- Lee HS, Park JH, Kim SJ, Kwon SJ, Kwon J. 2010. A cooperative activation loop among SWI/SNF, γ -H2AX and H3 acetylation for DNA double-strand break repair. *EMBO J* **29**: 1434–1445.
- Li Y, Bolderson E, Kumar R, Muniandy PA, Xue Y, Richard DJ, Seidman M, Pandita TK, Khanna KK, Wang W. 2009. HSSB1 and hSSB2 form similar multiprotein complexes that participate in DNA damage response. *J Biol Chem* **284**: 23525–23531.
- Li GY, McCulloch RD, Fenton AL, Cheung M, Meng L, Ikura M, Koch CA. 2010. Structure and identification of ADP-ribose recognition motifs of APLF and role in the DNA damage response. *Proc Natl Acad Sci* **107**: 9129–9134.
- Li L, Halaby MJ, Hakem A, Cardoso R, El Ghamrasni S, Harding S, Chan N, Bristow R, Sanchez O, Durocher D, et al. 2010. Rnf8 deficiency impairs class switch recombination, spermatogenesis, and genomic integrity and predisposes for cancer. *J Exp Med* **207**: 983–997.
- Li X, Corsa CA, Pan PW, Wu L, Ferguson D, Yu X, Min J, Dou Y. 2010. MOF and H4 K16 acetylation play important roles in DNA damage repair by modulating recruitment of DNA damage repair protein Mdc1. *Mol Cell Biol* **30**: 5335–5347.
- Lieber MR. 2010. The mechanism of double-strand DNA break repair by the nonhomologous DNA end-joining pathway. *Annu Rev Biochem* **79**: 181–211.
- Lieber MR, Wilson TE. 2010. SnapShot: Nonhomologous DNA end joining (NHEJ). *Cell* **142**: 496–496 e491.
- Lilley CE, Chaurushiya MS, Boutell C, Landry S, Suh J, Panier S, Everett RD, Stewart GS, Durocher D, Weitzman MD. 2010. A viral E3 ligase targets RNF8 and RNF168 to control histone ubiquitination and DNA damage responses. *EMBO J* **29**: 943–955.
- Limoli CL, Ward JF. 1993. A new method for introducing double-strand breaks into cellular DNA. *Radiat Res* **134**: 160–169.
- Lindahl T, Barnes DE. 2000. Repair of endogenous DNA damage. *Cold Spring Harb Symp Quant Biol* **65**: 127–133.
- Lindroos HB, Strom L, Itoh T, Katou Y, Shirahige K, Sjogren C. 2006. Chromosomal association of the Smc5/6 complex reveals that it functions in differently regulated pathways. *Mol Cell* **22**: 755–767.
- Lisby M, Mortensen UH, Rothstein R. 2003. Colocalization of multiple DNA double-strand breaks at a single Rad52 repair centre. *Nat Cell Biol* **5**: 572–577.
- Lisby M, Barlow JH, Burgess RC, Rothstein R. 2004. Choreography of the DNA damage response: spatiotemporal relationships among checkpoint and repair proteins. *Cell* **118**: 699–713.
- Liu Y, Subrahmanyam R, Chakraborty T, Sen R, Desiderio S. 2007. A plant homeodomain in RAG-2 that binds hypermethylated lysine 4 of histone H3 is necessary for efficient antigen-receptor-gene rearrangement. *Immunity* **27**: 561–571.
- Liu J, Doty T, Gibson B, Heyer WD. 2010. Human BRCA2 protein promotes RAD51 filament formation on RPA-covered single-stranded DNA. *Nat Struct Mol Biol* **17**: 1260–1262.
- Liu T, Ghosal G, Yuan J, Chen J, Huang J. 2010. FAN1 acts with FANCI-FANCD2 to promote DNA interstrand cross-link repair. *Science* **329**: 693–696.
- Lloyd J, Chapman JR, Clapperton JA, Haire LF, Hartsuiker E, Li J, Carr AM, Jackson SP, Smerdon SJ. 2009. A supramodular FHA/BRCT-repeat architecture mediates Nbs1 adaptor function in response to DNA damage. *Cell* **139**: 100–111.
- Longhese MP, Bonetti D, Guerini I, Manfrini N, Clerici M. 2009. DNA double-strand breaks in meiosis: checking their formation, processing and repair. *DNA Repair (Amst)* **8**: 1127–1138.
- Longhese MP, Bonetti D, Manfrini N, Clerici M. 2010. Mechanisms and regulation of DNA end resection. *EMBO J* **29**: 2864–2874.
- Lou Z, Minter-Dykhouse K, Franco S, Gostissa M, Rivera MA, Celeste A, Manis JP, van Deursen J, Nussenzweig A, Paull TT, et al. 2006. MDC1 maintains genomic stability by participating in the amplification of ATM-dependent DNA damage signals. *Mol Cell* **21**: 187–200.
- Luger K, Mader AW, Richmond RK, Sargent DE, Richmond TJ. 1997. Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature* **389**: 251–260.
- Luijsterburg MS, Dinant C, Lans H, Stap J, Wiernasz E, Lagerwerf S, Warmerdam DO, Lindh M, Brink MC, Dobrucki JW, et al. 2009. Heterochromatin protein 1 is recruited to various types of DNA damage. *J Cell Biol* **185**: 577–586.
- Lukas C, Falck J, Bartkova J, Bartek J, Lukas J. 2003. Distinct spatiotemporal dynamics of mammalian checkpoint regulators induced by DNA damage. *Nat Cell Biol* **5**: 255–260.
- Lukas C, Bartek J, Lukas J. 2005. Imaging of protein movement induced by chromosomal breakage: tiny ‘local’ lesions pose great ‘global’ challenges. *Chromosoma* **114**: 146–154.

- MacKay C, Declais AC, Lundin C, Agostinho A, Deans AJ, MacArtney TJ, Hofmann K, Gartner A, West SC, Helleday T, et al. 2010. Identification of KIAA1018/FAN1, a DNA repair nuclease recruited to DNA damage by monoubiquitinated FANCD2. *Cell* **142**: 65–76.
- Macurek L, Lindqvist A, Voets O, Kool J, Vos HR, Medema RH. 2010. Wip1 phosphatase is associated with chromatin and dephosphorylates γ H2AX to promote checkpoint inhibition. *Oncogene* **29**: 2281–2291.
- Mah LJ, El-Osta A, Karagiannis TC. 2010. γ H2AX as a molecular marker of aging and disease. *Epigenetics* **5**: 129–136.
- Mailand N, Bekker-Jensen S, Fastrup H, Melander F, Bartek J, Lukas C, Lukas J. 2007. RNF8 ubiquitylates histones at DNA double-strand breaks and promotes assembly of repair proteins. *Cell* **131**: 887–900.
- Manke IA, Lowery DM, Nguyen A, Yaffe MB. 2003. BRCT repeats as phosphopeptide-binding modules involved in protein targeting. *Science* **302**: 636–639.
- Mari PO, Florea BI, Persengiev SP, Verkaik NS, Bruggenwirth HT, Modesti M, Giglia-Mari G, Bezstarosti K, Demmers JA, Luidert TM, et al. 2006. Dynamic assembly of end-joining complexes requires interaction between Ku70/80 and XRCC4. *Proc Natl Acad Sci* **103**: 18597–18602.
- Maser RS, Monsen KJ, Nelms BE, Petrini JH. 1997. hMre11 and hRad50 nuclear foci are induced during the normal cellular response to DNA double-strand breaks. *Mol Cell Biol* **17**: 6087–6096.
- Massip L, Caron P, Iacovoni JS, Trouche D, Legube G. 2010. Deciphering the chromatin landscape induced around DNA double strand breaks. *Cell Cycle* **9**: 2963–2972.
- Masutani M, Nozaki T, Nishiyama E, Shimokawa T, Tachi Y, Suzuki H, Nakagama H, Wakabayashi K, Sugimura T. 1999. Function of poly(ADP-ribose) polymerase in response to DNA damage: gene-disruption study in mice. *Mol Cell Biochem* **193**: 149–152.
- Matsuoka S, Ballif BA, Smogorzewska A, McDonald ER III, Hurov KE, Luo J, Bakalarski CE, Zhao Z, Solimini N, Lerenthal Y, et al. 2007. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* **316**: 1160–1166.
- Matsuzaki K, Shinohara A, Shinohara M. 2008. Forkhead-associated domain of yeast Xrs2, a homolog of human Nbs1, promotes nonhomologous end joining through interaction with a ligase IV partner protein, Lif1. *Genetics* **179**: 213–225.
- Matthews AG, Kuo AJ, Ramon-Maiques S, Han S, Champagne KS, Ivanov D, Gallardo M, Carney D, Cheung P, Ciccone DN, et al. 2007. RAG2 PHD finger couples histone H3 lysine 4 trimethylation with V(D)J recombination. *Nature* **450**: 1106–1110.
- Mazon G, Mimitou EP, Symington LS. 2010. SnapShot: Homologous recombination in DNA double-strand break repair. *Cell* **142**: 646, 646 e641.
- McBryant SJ, Adams VH, Hansen JC. 2006. Chromatin architectural proteins. *Chromosome Res* **14**: 39–51.
- McVey M, Lee SE. 2008. MMEJ repair of double-strand breaks (director's cut): deleted sequences and alternative endings. *Trends Genet* **24**: 529–538.
- Meier A, Fiegler H, Munoz P, Ellis P, Rigler D, Langford C, Blasco MA, Carter N, Jackson SP. 2007. Spreading of mammalian DNA-damage response factors studied by ChIP-chip at damaged telomeres. *EMBO J* **26**: 2707–2718.
- Melander F, Bekker-Jensen S, Falck J, Bartek J, Mailand N, Lukas J. 2008. Phosphorylation of SDT repeats in the MDC1 N terminus triggers retention of NBS1 at the DNA damage-modified chromatin. *J Cell Biol* **181**: 213–226.
- Menissier-de Murcia J, Molinete M, Gradwohl G, Simonin F, de Murcia G. 1989. Zinc-binding domain of poly(ADP-ribose) polymerase participates in the recognition of single strand breaks on DNA. *J Mol Biol* **210**: 229–233.
- Menissier de Murcia J, Ricoul M, Tartier L, Niedergang C, Huber A, Dantzer F, Schreiber V, Ame JC, Dierich A, LeMeur M, et al. 2003. Functional interaction between PARP-1 and PARP-2 in chromosome stability and embryonic development in mouse. *EMBO J* **22**: 2255–2263.
- Merkle D, Douglas P, Moorhead GB, Leonenko Z, Yu Y, Cramb D, Bazett-Jones DP, Lees-Miller SP. 2002. The DNA-dependent protein kinase interacts with DNA to form a protein-DNA complex that is disrupted by phosphorylation. *Biochemistry* **41**: 12706–12714.
- Messick TE, Greenberg RA. 2009. The ubiquitin landscape at DNA double-strand breaks. *J Cell Biol* **187**: 319–326.
- Miller KM, Tjeertes JV, Coates J, Legube G, Polo SE, Britton S, Jackson SP. 2010. Human HDAC1 and HDAC2 function in the DNA-damage response to promote DNA nonhomologous end-joining. *Nat Struct Mol Biol* **17**: 1144–1151.
- Mimitou EP, Symington LS. 2009. Nucleases and helicases take center stage in homologous recombination. *Trends Biochem Sci* **34**: 264–272.
- Mimori T, Hardin JA. 1986. Mechanism of interaction between Ku protein and DNA. *J Biol Chem* **261**: 10375–10379.
- Misteli T, Soutoglou E. 2009. The emerging role of nuclear architecture in DNA repair and genome maintenance. *Nat Rev Mol Cell Biol* **10**: 243–254.
- Moggs JG, Grandi P, Quivy JP, Jonsson ZO, Hubscher U, Becker PB, Almouzni G. 2000. A CAF-1-PCNA-mediated chromatin assembly pathway triggered by sensing DNA damage. *Mol Cell Biol* **20**: 1206–1218.
- Mohammad DH, Yaffe MB. 2009. 14-3-3 proteins, FHA domains and BRCT domains in the DNA damage response. *DNA Repair (Amst)* **8**: 1009–1017.
- Moldovan GL, D'Andrea AD. 2009. How the fanconi anemia pathway guards the genome. *Annu Rev Genet* **43**: 223–249.
- Moon SH, Lin L, Zhang X, Nguyen TA, Darlington Y, Waldman AS, Lu X, Donehower LA. 2010. Wild-type p53-induced phosphatase 1 dephosphorylates histone variant γ H2AX and suppresses DNA double strand break repair. *J Biol Chem* **285**: 12935–12947.
- Moreno-Herrero F, de Jager M, Dekker NH, Kanaar R, Wyman C, Dekker C. 2005. Mesoscale conformational changes in the DNA-repair complex Rad50/Mre11/Nbs1 upon binding DNA. *Nature* **437**: 440–443.
- Morris JR, Solomon E. 2004. BRCA1: BARD1 induces the formation of conjugated ubiquitin structures, dependent on K6 of ubiquitin, in cells during DNA replication and repair. *Hum Mol Genet* **13**: 807–817.
- Morris JR, Boutell C, Keppler M, Densham R, Weekes D, Alamshah A, Butler L, Galanty Y, Pangon L, Kiuchi T, et al. 2009. The SUMO modification pathway is involved in the BRCA1 response to genotoxic stress. *Nature* **462**: 886–890.
- Morrison AJ, Highland J, Krogan NJ, Arbel-Eden A, Greenblatt JF, Haber JE, Shen X. 2004. INO80 and γ H2AX interaction links ATP-dependent chromatin remodeling to DNA damage repair. *Cell* **119**: 767–775.
- Mortusewicz O, Schermelleh L, Walter J, Cardoso MC, Leonhardt H. 2005. Recruitment of DNA methyltransferase I to DNA repair sites. *Proc Natl Acad Sci* **102**: 8905–8909.
- Mortusewicz O, Ame JC, Schreiber V, Leonhardt H. 2007. Feedback-regulated poly(ADP-ribosylation) by PARP-1 is required for rapid response to DNA damage in living cells. *Nucleic Acids Res* **35**: 7665–7675.

Polo and Jackson

- Mortusewicz O, Leonhardt H, Cardoso MC. 2008. Spatiotemporal dynamics of regulatory protein recruitment at DNA damage sites. *J Cell Biochem* **104**: 1562–1569.
- Moynahan ME, Jasin M. 2010. Mitotic homologous recombination maintains genomic stability and suppresses tumorigenesis. *Nat Rev Mol Cell Biol* **11**: 196–207.
- Murr R, Loizou JI, Yang YG, Cuenin C, Li H, Wang ZQ, Herceg Z. 2006. Histone acetylation by Trapp-Tip60 modulates loading of repair proteins and repair of DNA double-strand breaks. *Nat Cell Biol* **8**: 91–99.
- Nagai S, Dubrana K, Tsai-Pflugfelder M, Davidson MB, Roberts TM, Brown GW, Varela E, Hediger F, Gasser SM, Krogan NJ. 2008. Functional targeting of DNA damage to a nuclear pore-associated SUMO-dependent ubiquitin ligase. *Science* **322**: 597–602.
- Nagy Z, Soutoglou E. 2009. DNA repair: easy to visualize, difficult to elucidate. *Trends Cell Biol* **19**: 617–629.
- Nakada D, Matsumoto K, Sugimoto K. 2003. ATM-related Tel1 associates with double-strand breaks through an Xrs2-dependent mechanism. *Genes Dev* **17**: 1957–1962.
- Nakada S, Chen GI, Gingras AC, Durocher D. 2008. PP4 is a γ H2AX phosphatase required for recovery from the DNA damage checkpoint. *EMBO Rep* **9**: 1019–1026.
- Nakada S, Tai I, Panier S, Al-Hakim A, Iemura S, Juang YC, O'Donnell L, Kumakubo A, Munro M, Sicheri F, et al. 2010. Non-canonical inhibition of DNA damage-dependent ubiquitination by OTUB1. *Nature* **466**: 941–946.
- Nakamura TM, Du LL, Redon C, Russell P. 2004. Histone H2A phosphorylation controls Crb2 recruitment at DNA breaks, maintains checkpoint arrest, and influences DNA repair in fission yeast. *Mol Cell Biol* **24**: 6215–6230.
- Nakamura AJ, Rao VA, Pommier Y, Bonner WM. 2010. The complexity of phosphorylated H2AX foci formation and DNA repair assembly at DNA double-strand breaks. *Cell Cycle* **9**: 389–397.
- Nazarov IB, Smirnova AN, Krutilina RI, Svetlova MP, Solovjeva LV, Nikiforov AA, Oei SL, Zalenskaya IA, Yau PM, Bradbury EM, et al. 2003. Dephosphorylation of histone γ -H2AX during repair of DNA double-strand breaks in mammalian cells and its inhibition by calyculin A. *Radiat Res* **160**: 309–317.
- Nelms BE, Maser RS, MacKay JF, Lagally MG, Petrini JH. 1998. In situ visualization of DNA double-strand break repair in human fibroblasts. *Science* **280**: 590–592.
- Nelson G, Buhmann M, von Zglinicki T. 2009. DNA damage foci in mitosis are devoid of 53BP1. *Cell Cycle* **8**: 3379–3383.
- Nicassio F, Corrado N, Vissers JH, Areces LB, Bergink S, Marteijn JA, Geverts B, Houtsmuller AB, Vermeulen W, Di Fiore PP, et al. 2007. Human USP3 is a chromatin modifier required for S phase progression and genome stability. *Curr Biol* **17**: 1972–1977.
- Nick McElhinny SA, Snowden CM, McCarville J, Ramsden DA. 2000. Ku recruits the XRCC4-ligase IV complex to DNA ends. *Mol Cell Biol* **20**: 2996–3003.
- Nijman SM, Huang TT, Dirac AM, Brummelkamp TR, Kerkhoven RM, D'Andrea AD, Bernards R. 2005. The deubiquitinating enzyme USP1 regulates the Fanconi anemia pathway. *Mol Cell Biol* **25**: 331–339.
- Noon AT, Shibata A, Rief N, Loblrich M, Stewart GS, Jeggo PA, Goodarzi AA. 2010. 53BP1-dependent robust localized KAP-1 phosphorylation is essential for heterochromatic DNA double-strand break repair. *Nat Cell Biol* **12**: 177–184.
- Oberdoerffer P, Michan S, McVay M, Mostoslavsky R, Vann J, Park SK, Hartlerode A, Stegmuller J, Hafner A, Loerch P, et al. 2008. SIRT1 redistribution on chromatin promotes genomic stability but alters gene expression during aging. *Cell* **135**: 907–918.
- Oda H, Hubner MR, Beck DB, Vermeulen M, Hurwitz J, Spector DL, Reinberg D. 2010. Regulation of the histone H4 monomethylase PR-Set7 by CRL4(Cdt2)-mediated PCNA-dependent degradation during DNA damage. *Mol Cell* **40**: 364–376.
- O'Hagan HM, Mohammad HP, Baylin SB. 2008. Double strand breaks can initiate gene silencing and SIRT1-dependent onset of DNA methylation in an exogenous promoter CpG island. *PLoS Genet* **4**: e1000155. doi: 10.1371/journal.pgen.1000155.
- Ohgushi H, Yoshihara K, Kamiya T. 1980. Bovine thymus poly(adenosine diphosphate ribose) polymerase. Physical properties and binding to DNA. *J Biol Chem* **255**: 6205–6211.
- Okano S, Lan L, Caldecott KW, Mori T, Yasui A. 2003. Spatial and temporal cellular responses to single-strand breaks in human cells. *Mol Cell Biol* **23**: 3974–3981.
- Palmbo PL, Wu D, Daley JM, Wilson TE. 2008. Recruitment of *Saccharomyces cerevisiae* Dnl4-Lif1 complex to a double-strand break requires interactions with Yku80 and the Xrs2 FHA domain. *Genetics* **180**: 1809–1819.
- Papamichos-Chronakis M, Krebs JE, Peterson CL. 2006. Interplay between I ν o80 and Swr1 chromatin remodeling enzymes regulates cell cycle checkpoint adaptation in response to DNA damage. *Genes Dev* **20**: 2437–2449.
- Pardo B, Gomez-Gonzalez B, Aguilera A. 2009. DNA repair in mammalian cells: DNA double-strand break repair: how to fix a broken relationship. *Cell Mol Life Sci* **66**: 1039–1056.
- Park JH, Park EJ, Lee HS, Kim SJ, Hur SK, Imbalzano AN, Kwon J. 2006. Mammalian SWI/SNF complexes facilitate DNA double-strand break repair by promoting γ -H2AX induction. *EMBO J* **25**: 3986–3997.
- Parrilla-Castellar ER, Arlander SJ, Karnitz L. 2004. Dial 9-1-1 for DNA damage: the Rad9-Hus1-Rad1 (9-1-1) clamp complex. *DNA Repair (Amst)* **3**: 1009–1014.
- Pataky K, Villanueva G, Liani A, Zgheib O, Jenkins N, Halazonetis DJ, Halazonetis TD, Brugger J. 2009. Microcollimator for micrometer-wide stripe irradiation of cells using 20–30 keV X rays. *Radiat Res* **172**: 252–259.
- Peng G, Yim EK, Dai H, Jackson AP, Burgt I, Pan MR, Hu R, Li K, Lin SY. 2009. BRIT1/MCPH1 links chromatin remodeling to DNA damage response. *Nat Cell Biol* **11**: 865–872.
- Petersen S, Casellas R, Reina-San-Martin B, Chen HT, Difilippantonio MJ, Wilson PC, Hanitsch L, Celeste A, Muramatsu M, Pilch DR, et al. 2001. AID is required to initiate Nbs1/ γ -H2AX focus formation and mutations at sites of class switching. *Nature* **414**: 660–665.
- Pickart CM. 2001. Mechanisms underlying ubiquitination. *Annu Rev Biochem* **70**: 503–533.
- Pinato S, Scandiuzzi C, Arnaudo N, Citterio E, Gaudino G, Penengo L. 2009. RNF168, a new RING finger, MIU-containing protein that modifies chromatin by ubiquitination of histones H2A and H2AX. *BMC Mol Biol* **10**: 55. doi: 10.1086/1471-2199-10-55.
- Pinato S, Gatti M, Scandiuzzi C, Confalonieri S, Penengo L. 2011. UMI, a novel RNF168 ubiquitin binding domain involved in the DNA damage signaling pathway. *Mol Cell Biol* **31**: 118–126.
- Pleschke JM, Kleczkowska HE, Strohm M, Althaus FR. 2000. Poly(ADP-ribose) binds to specific domains in DNA damage checkpoint proteins. *J Biol Chem* **275**: 40974–40980.
- Polanowska J, Martin JS, Garcia-Muse T, Petalcorin MI, Boulton SJ. 2006. A conserved pathway to activate BRCA1-dependent ubiquitylation at DNA damage sites. *EMBO J* **25**: 2178–2188.
- Polo SE, Roche D, Almouzni G. 2006. New histone incorporation marks sites of UV repair in human cells. *Cell* **127**: 481–493.
- Polo SE, Kaidi A, Baskcomb L, Galanty Y, Jackson SP. 2010. Regulation of DNA-damage responses and cell-cycle

- progression by the chromatin remodelling factor CHD4. *EMBO J* **29**: 3130–3139.
- Qin S, Parthun MR. 2006. Recruitment of the type B histone acetyltransferase Hat1p to chromatin is linked to DNA double-strand breaks. *Mol Cell Biol* **26**: 3649–3658.
- Richard DJ, Bolderson E, Cubeddu L, Wadsworth RI, Savage K, Sharma GG, Nicolette ML, Tsvetanov S, McIlwraith MJ, Pandita RK, et al. 2008. Single-stranded DNA-binding protein hSSB1 is critical for genomic stability. *Nature* **453**: 677–681.
- Richard DJ, Savage K, Bolderson E, Cubeddu L, So S, Ghita M, Chen DJ, White MF, Richard K, Prise KM et al. 2010. hSSB1 rapidly binds at the sites of DNA double-strand breaks and is required for the efficient recruitment of the MRN complex. *Nucleic Acids Res*. doi: 10.1093/nar/gkq1098.
- Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. 1998. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem* **273**: 5858–5868.
- Rogakou EP, Boon C, Redon C, Bonner WM. 1999. Megabase chromatin domains involved in DNA double-strand breaks in vivo. *J Cell Biol* **146**: 905–916.
- Rouse J, Jackson SP. 2002. Lcd1p recruits Mec1p to DNA lesions in vitro and in vivo. *Mol Cell* **9**: 857–869.
- Rozenzhak S, Mejia-Ramirez E, Williams JS, Schaffer L, Hammond JA, Head SR, Russell P. 2010. Rad3 decorates critical chromosomal domains with γ H2A to protect genome integrity during S-Phase in fission yeast. *PLoS Genet* **6**: e1001032. doi: 10.1371/journal.pgen.1001032.
- Rulten SL, Cortes-Ledesma F, Guo L, Iles NJ, Caldecott KW. 2008. APLF (C2orf13) is a novel component of poly(ADP-ribose) signaling in mammalian cells. *Mol Cell Biol* **28**: 4620–4628.
- Rupnik A, Lowndes NF, Grenon M. 2010. MRN and the race to the break. *Chromosoma* **119**: 115–135.
- Sacher M, Pfander B, Hoegge C, Jentsch S. 2006. Control of Rad52 recombination activity by double-strand break-induced SUMO modification. *Nat Cell Biol* **8**: 1284–1290.
- Sanders SL, Portoso M, Mata J, Bahler J, Allshire RC, Kouzarides T. 2004. Methylation of histone H4 lysine 20 controls recruitment of Crb2 to sites of DNA damage. *Cell* **119**: 603–614.
- Sanders SL, Arida AR, Phan FP. 2010. Requirement for the phospho-H2AX binding module of Crb2 in double-strand break targeting and checkpoint activation. *Mol Cell Biol* **30**: 4722–4731.
- San Filippo J, Sung P, Klein H. 2008. Mechanism of eukaryotic homologous recombination. *Annu Rev Biochem* **77**: 229–257.
- Santos MA, Huen MS, Jankovic M, Chen HT, Lopez-Contreras AJ, Klein IA, Wong N, Barbancho JL, Fernandez-Capetillo O, Nussenzweig MC, et al. 2010. Class switching and meiotic defects in mice lacking the E3 ubiquitin ligase RNF8. *J Exp Med* **207**: 973–981.
- Sartori AA, Lukas C, Coates J, Mistrik M, Fu S, Bartek J, Baer R, Lukas J, Jackson SP. 2007. Human CtIP promotes DNA end resection. *Nature* **450**: 509–514.
- Sato Y, Yoshikawa A, Mimura H, Yamashita M, Yamagata A, Fukai S. 2009. Structural basis for specific recognition of Lys 63-linked polyubiquitin chains by tandem UIMs of RAP80. *EMBO J* **28**: 2461–2468.
- Schermlle L, Carlton PM, Haese S, Shao L, Winoto L, Kner P, Burke B, Cardoso MC, Agard DA, Gustafsson MG, et al. 2008. Subdiffraction multicolor imaging of the nuclear periphery with 3D structured illumination microscopy. *Science* **320**: 1332–1336.
- Scully R, Chen J, Ochs RL, Keegan K, Hoekstra M, Feunteun J, Livingston DM. 1997. Dynamic changes of BRCA1 subnuclear location and phosphorylation state are initiated by DNA damage. *Cell* **90**: 425–435.
- Shanbhag NM, Rafalska-Metcalf IU, Balane-Bolivar C, Janicki SM, Greenberg RA. 2010. ATM-dependent chromatin changes silence transcription in cis to DNA double-strand breaks. *Cell* **141**: 970–981.
- Shao G, Lilli DR, Patterson-Fortin J, Coleman KA, Morrissey DE, Greenberg RA. 2009. The Rap80–BRCC36 de-ubiquitinating enzyme complex antagonizes RNF8–Ubc13-dependent ubiquitination events at DNA double strand breaks. *Proc Natl Acad Sci* **106**: 3166–3171.
- Sharma GG, So S, Gupta A, Kumar R, Cayrou C, Avvakumov N, Bhadra U, Pandita RK, Porteus MH, Chen DJ, et al. 2010. MOF and histone H4 acetylation at lysine 16 are critical for DNA damage response and double-strand break repair. *Mol Cell Biol* **30**: 3582–3595.
- Shim EY, Ma JL, Oum JH, Yanez Y, Lee SE. 2005. The yeast chromatin remodeler RSC complex facilitates end joining repair of DNA double-strand breaks. *Mol Cell Biol* **25**: 3934–3944.
- Shim EY, Hong SJ, Oum JH, Yanez Y, Zhang Y, Lee SE. 2007. RSC mobilizes nucleosomes to improve accessibility of repair machinery to the damaged chromatin. *Mol Cell Biol* **27**: 1602–1613.
- Shroff R, Arbel-Eden A, Pilch D, Ira G, Bonner WM, Petrini JH, Haber JE, Lichten M. 2004. Distribution and dynamics of chromatin modification induced by a defined DNA double-strand break. *Curr Biol* **14**: 1703–1711.
- Sims JJ, Cohen RE. 2009. Linkage-specific avidity defines the lysine 63-linked polyubiquitin-binding preference of rap80. *Mol Cell* **33**: 775–783.
- Singleton BK, Torres-Arzayus MI, Rottinghaus ST, Taccioli GE, Jeggo PA. 1999. The C terminus of Ku80 activates the DNA-dependent protein kinase catalytic subunit. *Mol Cell Biol* **19**: 3267–3277.
- Skaar JR, Richard DJ, Saraf A, Toschi A, Bolderson E, Florens L, Washburn MP, Khanna KK, Pagano M. 2009. INTS3 controls the hSSB1-mediated DNA damage response. *J Cell Biol* **187**: 25–32.
- Smeenk G, Wiegant WW, Vrolijk H, Solari AP, Pastink A, van Attikum H. 2010. The NuRD chromatin-remodeling complex regulates signaling and repair of DNA damage. *J Cell Biol* **190**: 741–749.
- Smits VA, Reaper PM, Jackson SP. 2006. Rapid PIKK-dependent release of Chk1 from chromatin promotes the DNA-damage checkpoint response. *Curr Biol* **16**: 150–159.
- Smogorzewska A, Desetty R, Saito TT, Schlabach M, Lach FP, Sowa ME, Clark AB, Kunkel TA, Harper JW, Colaiacovo MP, et al. 2010. A genetic screen identifies FAN1, a Fanconi anemia-associated nuclease necessary for DNA interstrand crosslink repair. *Mol Cell* **39**: 36–47.
- Sobhian B, Shao G, Lilli DR, Culhane AC, Moreau LA, Xia B, Livingston DM, Greenberg RA. 2007. RAP80 targets BRCA1 to specific ubiquitin structures at DNA damage sites. *Science* **316**: 1198–1202.
- Sofueva S, Du LL, Limbo O, Williams JS, Russell P. 2010. BRCT domain interactions with phospho-histone H2A target Crb2 to chromatin at double-strand breaks and maintain the DNA damage checkpoint. *Mol Cell Biol* **30**: 4732–4743.
- Soutoglou E, Misteli T. 2008. Activation of the cellular DNA damage response in the absence of DNA lesions. *Science* **320**: 1507–1510.
- Soutoglou E, Dorn JF, Sengupta K, Jasin M, Nussenzweig A, Ried T, Danuser G, Misteli T. 2007. Positional stability of single double-strand breaks in mammalian cells. *Nat Cell Biol* **9**: 675–682.

Polo and Jackson

- Spycher C, Miller ES, Townsend K, Pavic L, Morrice NA, Janscak P, Stewart GS, Stucki M. 2008. Constitutive phosphorylation of MDC1 physically links the MRE11–RAD50–NBS1 complex to damaged chromatin. *J Cell Biol* **181**: 227–240.
- Stap J, Krawczyk PM, Van Oven CH, Barendsen GW, Essers J, Kanaar R, Aten JA. 2008. Induction of linear tracks of DNA double-strand breaks by α -particle irradiation of cells. *Nat Methods* **5**: 261–266.
- Stewart GS, Panier S, Townsend K, Al-Hakim AK, Kolas NK, Miller ES, Nakada S, Ylanko J, Olivarius S, Mendez M, et al. 2009. The RIDDLE syndrome protein mediates a ubiquitin-dependent signaling cascade at sites of DNA damage. *Cell* **136**: 420–434.
- Stiff T, O'Driscoll M, Rief N, Iwabuchi K, Lohrich M, Jeggo PA. 2004. ATM and DNA-PK function redundantly to phosphorylate H2AX after exposure to ionizing radiation. *Cancer Res* **64**: 2390–2396.
- Stracker TH, Usui T, Petrini JH. 2009. Taking the time to make important decisions: the checkpoint effector kinases Chk1 and Chk2 and the DNA damage response. *DNA Repair (Amst)* **8**: 1047–1054.
- Strom L, Lindroos HB, Shirahige K, Sjogren C. 2004. Postreplicative recruitment of cohesin to double-strand breaks is required for DNA repair. *Mol Cell* **16**: 1003–1015.
- Stucki M, Clapperton JA, Mohammad D, Yaffe MB, Smerdon SJ, Jackson SP. 2005. MDC1 directly binds phosphorylated histone H2AX to regulate cellular responses to DNA double-strand breaks. *Cell* **123**: 1213–1226.
- Sugawara N, Haber JE. 2006. Repair of DNA double strand breaks: in vivo biochemistry. *Methods Enzymol* **408**: 416–429.
- Sun Y, Jiang X, Xu Y, Ayrapetov MK, Moreau LA, Whetstone JR, Price BD. 2009. Histone H3 methylation links DNA damage detection to activation of the tumour suppressor Tip60. *Nat Cell Biol* **11**: 1376–1382.
- Sy SM, Huen MS, Chen J. 2009. PALB2 is an integral component of the BRCA complex required for homologous recombination repair. *Proc Natl Acad Sci* **106**: 7155–7160.
- Szilard RK, Jacques PE, Laramée L, Cheng B, Galicia S, Bataille AR, Yeung M, Mendez M, Bergeron M, Robert F, et al. 2010. Systematic identification of fragile sites via genome-wide location analysis of γ -H2AX. *Nat Struct Mol Biol* **17**: 299–305.
- Takai H, Smogorzewska A, de Lange T. 2003. DNA damage foci at dysfunctional telomeres. *Curr Biol* **13**: 1549–1556.
- Tamburini BA, Tyler JK. 2005. Localized histone acetylation and deacetylation triggered by the homologous recombination pathway of double-strand DNA repair. *Mol Cell Biol* **25**: 4903–4913.
- Tartier L, Spenlehauer C, Newman HC, Folkard M, Prise KM, Michael BD, Menissier-de Murcia J, de Murcia G. 2003. Local DNA damage by proton microbeam irradiation induces poly(ADP-ribose) synthesis in mammalian cells. *Mutagenesis* **18**: 411–416.
- Teo SH, Jackson SP. 2000. Lif1p targets the DNA ligase Lig4p to sites of DNA double-strand breaks. *Curr Biol* **10**: 165–168.
- Thorslund T, McIlwraith MJ, Compton SA, Lekomtsev S, Petronczki M, Griffith JD, West SC. 2010. The breast cancer tumor suppressor BRCA2 promotes the specific targeting of RAD51 to single-stranded DNA. *Nat Struct Mol Biol* **17**: 1263–1265.
- Timinszky G, Till S, Hassa PO, Hothorn M, Kustatscher G, Nijmeijer B, Colombelli J, Altmeyer M, Stelzer EH, Scheffzek K, et al. 2009. A macrodomain-containing histone rearranges chromatin upon sensing PARP1 activation. *Nat Struct Mol Biol* **16**: 923–929.
- Torres-Rosell J, Sunjevaric I, De Piccoli G, Sacher M, Eckert-Boulet N, Reid R, Jentsch S, Rothstein R, Aragon L, Lisby M. 2007. The Smc5–Smc6 complex and SUMO modification of Rad52 regulates recombinational repair at the ribosomal gene locus. *Nat Cell Biol* **9**: 923–931.
- Tsai AG, Lieber MR. 2010. Mechanisms of chromosomal rearrangement in the human genome. *BMC Genomics* **11**: S1. doi: 10.1186/1471-2164-11-S1-S1.
- Tsukuda T, Fleming AB, Nickoloff JA, Osley MA. 2005. Chromatin remodelling at a DNA double-strand break site in *Saccharomyces cerevisiae*. *Nature* **438**: 379–383.
- Uematsu N, Weterings E, Yano K, Morotomi-Yano K, Jakob B, Taucher-Scholz G, Mari PO, van Gent DC, Chen BP, Chen DJ. 2007. Autophosphorylation of DNA-PKCS regulates its dynamics at DNA double-strand breaks. *J Cell Biol* **177**: 219–229.
- Unal E, Arbel-Eden A, Sattler U, Shroff R, Lichten M, Haber JE, Koshland D. 2004. DNA damage response pathway uses histone modification to assemble a double-strand break-specific cohesin domain. *Mol Cell* **16**: 991–1002.
- Unal E, Heidinger-Pauli JM, Koshland D. 2007. DNA double-strand breaks trigger genome-wide sister-chromatid cohesion through Eco1 (Ctf7). *Science* **317**: 245–248.
- Urnov FD, Miller JC, Lee YL, Beausejour CM, Rock JM, Augustus S, Jamieson AC, Porteus MH, Gregory PD, Holmes MC. 2005. Highly efficient endogenous human gene correction using designed zinc-finger nucleases. *Nature* **435**: 646–651.
- Uziel T, Lerenthal Y, Moyal L, Andegeko Y, Mittelman L, Shiloh Y. 2003. Requirement of the MRN complex for ATM activation by DNA damage. *EMBO J* **22**: 5612–5621.
- van Attikum H, Gasser SM. 2009. Crosstalk between histone modifications during the DNA damage response. *Trends Cell Biol* **19**: 207–217.
- van Attikum H, Fritsch O, Hohn B, Gasser SM. 2004. Recruitment of the INO80 complex by H2A phosphorylation links ATP-dependent chromatin remodeling with DNA double-strand break repair. *Cell* **119**: 777–788.
- van Attikum H, Fritsch O, Gasser SM. 2007. Distinct roles for SWR1 and INO80 chromatin remodeling complexes at chromosomal double-strand breaks. *EMBO J* **26**: 4113–4125.
- van Veelen LR, Cervelli T, van de Rakt MW, Theil AF, Essers J, Kanaar R. 2005. Analysis of ionizing radiation-induced foci of DNA damage repair proteins. *Mutat Res* **574**: 22–33.
- van Vugt MA, Gardino AK, Lindling R, Ostheimer GJ, Reinhardt HC, Ong SE, Tan CS, Miao H, Keezer SM, Li J, et al. 2010. A mitotic phosphorylation feedback network connects Cdk1, Plk1, 53BP1, and Chk2 to inactivate the G2/M DNA damage checkpoint. *PLoS Biol* **8**: e1000287. doi: 10.1371/journal.pbio.1000287.
- Verdun RE, Crabbe L, Haggblom C, Karlseder J. 2005. Functional human telomeres are recognized as DNA damage in G2 of the cell cycle. *Mol Cell* **20**: 551–561.
- Walker JR, Corpina RA, Goldberg J. 2001. Structure of the Ku heterodimer bound to DNA and its implications for double-strand break repair. *Nature* **412**: 607–614.
- Wang B, Elledge SJ. 2007. Ubc13/Rnf8 ubiquitin ligases control foci formation of the Rap80/Abraxas/Brc1/Brc36 complex in response to DNA damage. *Proc Natl Acad Sci* **104**: 20759–20763.
- Wang ZQ, Auer B, Stingl L, Berghammer H, Haidacher D, Schweiger M, Wagner EF. 1995. Mice lacking ADPRT and poly(ADP-ribosyl)ation develop normally but are susceptible to skin disease. *Genes Dev* **9**: 509–520.
- Wang M, Wu W, Rosidi B, Zhang L, Wang H, Iliakis G. 2006. PARP-1 and Ku compete for repair of DNA double strand

- breaks by distinct NHEJ pathways. *Nucleic Acids Res* **34**: 6170–6182.
- Ward IM, Chen J. 2001. Histone H2AX is phosphorylated in an ATR-dependent manner in response to replicational stress. *J Biol Chem* **276**: 47759–47762.
- Ward IM, Reina-San-Martin B, Oлару A, Minn K, Tamada K, Lau JS, Cascalho M, Chen L, Nussenzweig A, Livak F, et al. 2004. 53BP1 is required for class switch recombination. *J Cell Biol* **165**: 459–464.
- Watanabe K, Iwabuchi K, Sun J, Tsuji Y, Tani T, Tokunaga K, Date T, Hashimoto M, Yamaizumi M, Tateishi S. 2009. RAD18 promotes DNA double-strand break repair during G1 phase through chromatin retention of 53BP1. *Nucleic Acids Res* **37**: 2176–2193.
- Wellinger RJ. 2010. When the caps fall off: responses to telomere uncapping in yeast. *FEBS Lett* **584**: 3734–3740.
- Williams RS, Moncalian G, Williams JS, Yamada Y, Limbo O, Shin DS, Grocock LM, Cahill D, Hitomi C, Guenther G, et al. 2008. Mre11 dimers coordinate DNA end bridging and nuclease processing in double-strand-break repair. *Cell* **135**: 97–109.
- Williams RS, Dodson GE, Limbo O, Yamada Y, Williams JS, Guenther G, Classen S, Glover JN, Iwasaki H, Russell P, et al. 2009. Nbs1 flexibly tethers Ctp1 and Mre11–Rad50 to coordinate DNA double-strand break processing and repair. *Cell* **139**: 87–99.
- Wold MS. 1997. Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu Rev Biochem* **66**: 61–92.
- Wood JL, Singh N, Mer G, Chen J. 2007. MCPH1 functions in an H2AX-dependent but MDC1-independent pathway in response to DNA damage. *J Biol Chem* **282**: 35416–35423.
- Wu L, Luo K, Lou Z, Chen J. 2008. MDC1 regulates intra-S-phase checkpoint by targeting NBS1 to DNA double-strand breaks. *Proc Natl Acad Sci* **105**: 11200–11205.
- Wu J, Huen MS, Lu LY, Ye L, Dou Y, Ljungman M, Chen J, Yu X. 2009a. Histone ubiquitination associates with BRCA1-dependent DNA damage response. *Mol Cell Biol* **29**: 849–860.
- Wu J, Prindle MJ, Dressler GR, Yu X. 2009b. PTIP regulates 53BP1 and SMC1 at the DNA damage sites. *J Biol Chem* **284**: 18078–18084.
- Xia B, Sheng Q, Nakanishi K, Ohashi A, Wu J, Christ N, Liu X, Jasin M, Couch FJ, Livingston DM. 2006. Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2. *Mol Cell* **22**: 719–729.
- Xiao A, Li H, Shechter D, Ahn SH, Fabrizio LA, Erdjument-Bromage H, Ishibe-Murakami S, Wang B, Tempst P, Hofmann K, et al. 2009. WSTF regulates the H2A.X DNA damage response via a novel tyrosine kinase activity. *Nature* **457**: 57–62.
- Xie A, Puget N, Shim I, Odate S, Jarzyna I, Bassing CH, Alt FW, Scully R. 2004. Control of sister chromatid recombination by histone H2AX. *Mol Cell* **16**: 1017–1025.
- Xie A, Kwok A, Scully R. 2009. Role of mammalian Mre11 in classical and alternative nonhomologous end joining. *Nat Struct Mol Biol* **16**: 814–818.
- Xu Y, Sun Y, Jiang X, Ayrapetov MK, Moskwa P, Yang S, Weinstock DM, Price BD. 2010. The p400 ATPase regulates nucleosome stability and chromatin ubiquitination during DNA repair. *J Cell Biol* **191**: 31–43.
- Yan Q, Dutt S, Xu R, Graves K, Juszczynski P, Manis JP, Shipp MA. 2009. BBAP monoubiquitylates histone H4 at lysine 91 and selectively modulates the DNA damage response. *Mol Cell* **36**: 110–120.
- Yano K, Morotomi-Yano K, Wang SY, Uematsu N, Lee KJ, Asaithamby A, Weterings E, Chen DJ. 2008. Ku recruits XLF to DNA double-strand breaks. *EMBO Rep* **9**: 91–96.
- Yin B, Savic V, Juntilla MM, Bredemeyer AL, Yang-Iott KS, Helmink BA, Koretzky GA, Sleckman BP, Bassing CH. 2009. Histone H2AX stabilizes broken DNA strands to suppress chromosome breaks and translocations during V(D)J recombination. *J Exp Med* **206**: 2625–2639.
- You Z, Bailis JM. 2010. DNA damage and decisions: CtIP coordinates DNA repair and cell cycle checkpoints. *Trends Cell Biol* **20**: 402–409.
- You Z, Chahwan C, Bailis J, Hunter T, Russell P. 2005. ATM activation and its recruitment to damaged DNA require binding to the C terminus of Nbs1. *Mol Cell Biol* **25**: 5363–5379.
- Yu X, Fu S, Lai M, Baer R, Chen J. 2006. BRCA1 ubiquitinates its phosphorylation-dependent binding partner CtIP. *Genes Dev* **20**: 1721–1726.
- Yuan J, Chen J. 2010. MRE11–RAD50–NBS1 complex dictates DNA repair independent of H2AX. *J Biol Chem* **285**: 1097–1104.
- Yuan J, Adamski R, Chen J. 2010. Focus on histone variant H2AX: to be or not to be. *FEBS Lett* **584**: 3717–3724.
- Yurchenko V, Xue Z, Sadofsky MJ. 2006. SUMO modification of human XRCC4 regulates its localization and function in DNA double-strand break repair. *Mol Cell Biol* **26**: 1786–1794.
- Zarebski M, Wiernasz E, Dobrucki JW. 2009. Recruitment of heterochromatin protein 1 to DNA repair sites. *Cytometry A* **75**: 619–625.
- Zhang D, Zaugg K, Mak TW, Elledge SJ. 2006. A role for the deubiquitinating enzyme USP28 in control of the DNA-damage response. *Cell* **126**: 529–542.
- Zhang Y, Hefferin ML, Chen L, Shim EY, Tseng HM, Kwon Y, Sung P, Lee SE, Tomkinson AE. 2007. Role of Dnl4–Lif1 in nonhomologous end-joining repair complex assembly and suppression of homologous recombination. *Nat Struct Mol Biol* **14**: 639–646.
- Zhang F, Fan Q, Ren K, Andreassen PR. 2009a. PALB2 functionally connects the breast cancer susceptibility proteins BRCA1 and BRCA2. *Mol Cancer Res* **7**: 1110–1118.
- Zhang F, Ma J, Wu J, Ye L, Cai H, Xia B, Yu X. 2009b. PALB2 links BRCA1 and BRCA2 in the DNA-damage response. *Curr Biol* **19**: 524–529.
- Zhao X, Blobel G. 2005. A SUMO ligase is part of a nuclear multiprotein complex that affects DNA repair and chromosomal organization. *Proc Natl Acad Sci* **102**: 4777–4782.
- Zou L, Elledge SJ. 2003. Sensing DNA damage through ATRIP recognition of RPA–ssDNA complexes. *Science* **300**: 1542–1548.